

## Short Research Article

**The Solution of Homogeneous Liouville Fractional Differential Equations by Sumudu Transform Method****Abstract**

In this paper, we study the homogeneous Liouville fractional differential equations with constant coefficients. The solutions in terms of Mittag-Leffler of homogeneous Liouville fractional differential equations with constant coefficients are obtained by Sumudu transform method (STM). The results obtained by STM are illustrated by examples.

**Subject Classification:** 44A15, 44A99.

**Keywords:** Sumudu transform, Mittag-Leffler functions, Wright functions, Fractional differential equations.

## 1 Introduction

Many problems in physics, engineering and biology etc. are modeled via fractional differential equations such as diffusion, signal processing, electrochemistry, viscoelasticity [24, 27]. In literature numerous methods are available to solve fractional differential equations like power series method, iterative method, adomain decomposition method, transform method, monotone method etc. [1, 7, 9, 15, 18, 21, 26, 28]. Integral transform methods such as Fourier, Laplace, Mellin, and Hankel etc. were extensively used to study fractional differential equations [5, 6, 14, 16, 25]. In 1993, Watagulla [30, 31] introduced Sumudu transform and applied it to solve ordinary differential equations in control engineering problems. The complex inversion formula for Sumudu transform was proved by Weerakoon [28, 29] in 1994 and applied it to solve partial differential equations. Asiru studied the properties of Sumudu transform [1, 2, 3] and solved integral equations of convolution type [4] and discrete dynamical system [9]. Belgacem et al. [8] also established the properties of Sumudu transform. Kilicman et al. [19, 20] successfully applied Sumudu transform method to solve system of differential equations.

Kataetbeh and Belgacem [22, 24] obtained formulae for Sumudu transform of fractional derivatives such as Riemann-Liouville, Caputo and Miller-Ross using

Laplace- Sumudu duality property and obtained solutions of fractional differential equations. Bulut, Demiray and Tuluçe [10, 11, 12, 13, 17] have studied heat equations and wave equations by Sumudu transform method. In this paper we apply Sumudu transform method to obtain explicit solution of homogeneous fractional differential equations with constant coefficients. We also illustrate the STM by examples. section Preliminary Results, Notations and Terminology

In this section we give definitions and some basic results which are used in the next section.

**Definition 1.1** [24] A real function  $f(t)$ ,  $t \geq 0$  is said to be in space  $C_\mu$ ,  $\mu \in \mathbb{R}$  if there exists a real number  $n(> \mu)$ , such that  $f(t) = t^n f_1(t)$ , where  $f_1(t) \in C[0, \infty)$ , and is said to be in space  $C_\mu^k$  if and only if  $f^{(k)} \in C_\mu$ ,  $k \in \mathbb{N}$ .

**Definition 1.2** [21] The Liouville fractional integrals  $I_{0+}^\alpha f$  of order  $\alpha$  on the half-axis  $\mathbb{R}^+$  is defined as

$$(I_{0+}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \frac{f(t) dt}{(x-t)^{1-\alpha}}, \quad (x > 0; \Re(\alpha) > 0). \quad (1.1)$$

**Definition 1.3** [21] The Liouville fractional derivative  $D_{0+}^\alpha y$  of a function  $y(t)$  of order  $\alpha$  on the half-axis  $\mathbb{R}^+$  is given by

$$\begin{aligned} (D_{0+}^\alpha y)(t) &= \left(\frac{d}{dx}\right)^n ((I_{0+}^{n-\alpha} y)(x)) \\ &= \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dx}\right)^n \int_0^x \frac{y(t) dt}{(x-t)^{\alpha-n+1}}, \end{aligned} \quad (1.2)$$

with  $n = [\Re(\alpha)] + 1; \Re(\alpha) \geq 0; x > 0$ .

**Definition 1.4** [18] One parameter Mittag-Leffler function is denoted by  $E_\alpha(z)$  and is defined as

$$E_\alpha(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + 1)}, \quad \alpha \in \mathbb{C}, \Re(\alpha) > 0. \quad (1.3)$$

**Definition 1.5** [4] A two-parameter Mittag-Leffler function denoted by  $E_{\alpha,\beta}(z)$  and is defined as,

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}, \quad \alpha > 0, \beta > 0. \quad (1.4)$$

**Definition 1.6** [30] Consider a set  $A$  defined as

$$A = \{f(t) \mid \exists M, \tau_1, \tau_2 > 0, |f(t)| \leq Me^{\frac{|t|}{\tau_j}} \text{ if } t \in (-1)^j \times [0, \infty)\}. \quad (1.5)$$

For all real  $t \geq 0$ , the Sumudu transform of a function  $f(t) \in A$ , denoted by  $S[f(t)] = F(u)$ , is defined as

$$S[f(t)](u) = F(u) = \int_0^\infty e^{-t} f(ut) dt, \quad u \in (-\tau_1, \tau_2). \quad (1.6)$$

The function  $f(t)$  in equation (1.5) is called the inverse Sumudu transform of  $F(u)$  and is denoted by  $f(t) = S^{-1}[F(u)]$ .

**Definition 1.7** [22] The Sumudu transform of the Liouville fractional derivative (1.2) is given by

$$S[(D_{0+}^\alpha y)(t)](u) = u^{-\alpha} [Sy](u) - \sum_{j=1}^l d_j u^{-j} \quad (l-1 < \alpha \leq l; l \in \mathbb{N}) \quad (1.7)$$

where

$$d_j = (D_{0+}^{\alpha-j} y)(0+), \quad (j = 1, 2, \dots, l). \quad (1.8)$$

**Theorem 1.1** [8] Let  $F(u)$  and  $G(u)$  be the Sumudu transforms of  $f(t)$  and  $g(t)$  respectively. If

$$h(t) = (f * g)(t) = \int_0^t f(\tau)g(t - \tau) d\tau$$

where  $*$  denotes convolution of  $f$  and  $g$ , then the Sumudu transform of  $h(t)$  is

$$S[h(t)] = uF(u)G(u). \quad (1.9)$$

**Theorem 1.2** [7] Let  $n \geq 1$  and  $F(u)$  be the Sumudu transform of the function  $f(t)$ . The Sumudu transform of the  $n^{\text{th}}$  derivative of  $f(t)$ , denoted by  $S[f^{(n)}(t)](u) = F_n(u)$  is given by

$$\begin{aligned} S[f^{(n)}(t)](u) = F_n(u) &= \frac{F(u)}{u^n} - \frac{f(0)}{u^n} - \frac{f'(0)}{u^{(n-1)}} - \dots - \frac{f^{(n-1)}(0)}{u} \\ &= \frac{F(u)}{u^n} - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{u^{(n-k)}}. \end{aligned} \quad (1.10)$$

**Lemma 1.1** [14, 22] let  $\alpha, \beta, \lambda \in \mathbb{R}$  and  $\alpha > 0, \beta > 0, n \in \mathbb{N}$ . Then

$$S \left[ t^{\alpha n + \beta - 1} \left( \frac{\partial}{\partial \lambda} \right)^n E_{\alpha, \beta}(\lambda t^\alpha) \right] = \frac{n! u^{\alpha n + \beta - 1}}{(1 - \lambda u^\alpha)^{n+1}}. \quad (1.11)$$

In particular  $n = 0$  and  $\text{Re}(\frac{1}{u}) > \lambda$ , then

$$S[t^{\beta-1} E_{\alpha, \beta}(\lambda t^\alpha)](u) = \frac{u^{\beta-1}}{1 - \lambda u^\alpha}. \quad (1.12)$$

## 2 Homogenous Equations with Constant Coefficients :

In this section we apply Sumudu transform to obtain explicit solutions to the homogenous Liouville type fractional differential equation of the form

$$\sum_{k=0}^m A_k(D_{0+}^{\alpha_k}y)(t) + A_0y(t) = 0 \quad (t > 0; m \in \mathbb{N}; 0 < \alpha_1 < \dots < \alpha_m) \quad (2.1)$$

The conditions when solutions  $y_1(t), y_2(t), \dots, y_l(t), l \in \mathbb{N}$ , of equation (2.1) with  $l-1 < \alpha_m < l$  will be linearly independent and these solutions form the fundamental system of solutions given by

$$(D_{o+}^{\alpha-k}y_j)(0+) = \delta_{k,j} \quad (k, j = 1, 2, \dots, l), \quad (2.2)$$

where  $\delta_{k,j}$  is the Kronecker delta function.

**Theorem 2.1** *Let  $(t > 0; l - 1 < \alpha \leq l; l \in \mathbb{N}, \lambda \in \mathbb{R})$ . Then the functions*

$$y_j(t) = t^{\alpha-j} E_{\alpha, \alpha-j+1}(\lambda t^\alpha) \quad (j = 1, 2, \dots, l) \quad (2.3)$$

*yield the fundamental system of solution to the equation*

$$(D_{0+}^\alpha y)(t) - \lambda y(t) = 0. \quad (2.4)$$

**Proof :** Applying Sumudu transform on both sides of (2.4), we get

$$S[(D_{0+}^\alpha y)(t)](u) - \lambda S[y(t)](u) = 0. \quad (2.5)$$

Using (1.7), we get

$$\begin{aligned} u^{-\alpha}[Sy(t)](u) - \sum_{j=1}^l d_j u^{-j} - \lambda S[y(t)](u) &= 0 \\ (u^{-\alpha} - \lambda)S[y(t)](u) &= \sum_j^l d_j u^{-j} \\ S[y(t)](u) &= \sum_j^l d_j \frac{u^{-j}}{(u^{-\alpha} - \lambda)} \\ S[y(t)](u) &= \sum_j^l d_j \frac{u^{\alpha-j}}{(1 - \lambda u^\alpha)} \end{aligned} \quad (2.6)$$

Replacing  $\beta = \alpha - j + 1$ , in (1.12) and taking inverse Sumudu transform on both sides of (2.6), we get

$$y(t) = \sum_j^l d_j t^{\alpha-j} E_{\alpha, \alpha-j+1}(\lambda t^\alpha)$$

which gives the solution of the equation (2.4) as

$$y(t) = \sum_{j=1}^l d_j y_j(t),$$

where

$$y_j(t) = t^{\alpha-j} E_{\alpha, \alpha-j+1}(\lambda t^\alpha), j = 1, 2, \dots, l.$$

**Example 2.1** *The equation*

$$(D_{0+}^{l-\frac{1}{2}} y)(t) - \lambda y(t) = 0, \quad (t > 0; l \in \mathbb{N}; \lambda \in \mathbb{R}) \tag{2.7}$$

has its fundamental system of solution given by

$$y_j(t) = t^{l-j-\frac{1}{2}} E_{l-\frac{1}{2}, l-j+\frac{1}{2}}(\lambda t^{l-\frac{1}{2}}), \quad (j = 1, 2, \dots, l). \tag{2.8}$$

**Solution.** Applying Sumudu transform on both sides of (2.7), we get

$$S[(D_{0+}^{l-\frac{1}{2}} y)(t)](u) - \lambda S[y(t)](u) = 0. \tag{2.9}$$

Using (1.7), we get

$$u^{-(l-\frac{1}{2})} S[y(t)](u) - \sum_{j=1}^l d_j u^{-j} - \lambda S[y(t)](u) = 0$$

$$(u^{-(l-\frac{1}{2})} - \lambda)[Sy](u) = \sum_{j=1}^l d_j u^{-j}.$$

Using (2.2), we get

$$[Sy](u) = \sum_{j=1}^l d_j \frac{u^{l-j-\frac{1}{2}}}{(1 - \lambda u^{l-\frac{1}{2}})}. \tag{2.10}$$

Replacing  $\alpha = (l - \frac{1}{2})$  and  $\beta = l - j + \frac{1}{2}$  in (1.12) and taking inverse Sumudu transform on both sides of (2.10), we get the solution of (2.7) as

$$y(t) = \sum_{j=1}^l d_j y_j(t),$$

where

$$y_j(t) = t^{l-j-\frac{1}{2}} E_{l-\frac{1}{2}, l-j+\frac{1}{2}}(\lambda t^{l-\frac{1}{2}}), j = 1, 2, \dots, l.$$

### 3 Conclusion

We have obtained the solutions of homogeneous Liouville fractional differential equations with constant coefficients in terms of Mittag-Leffler by Sumudu transform method. The results obtained provides fundamental system of solutions of the considered problem. Results obtained are validated with some examples.

### References

- [1] Asiru, M.A. *Sumudu transform and the solution of integral equations of convolution type*, International Journal of Mathematical Education in Science and Technology, 32(6),906-910 (2001).
- [2] Asiru, M. A. *Further Properties of the Sumudu Transform and its Applications*, International Journal of Mathematical Education, Science and Technology, 33(2), 441-449 (2002).
- [3] Asiru, M.A. *Application of the Sumudu Transform to Discrete Dynamical Systems*, International Journal of Mathematical Education in Science and Technology, 34(6), 944-949 (2003).
- [4] Agarwal, R. P. *A propos d'unc note de M.Pierre Humbert*, C. R.Se'ances Acad. Sci., 236(21), 2031-2032 (1953).
- [5] Bodkhe. D. S and Panchal.S. K., *On Solution of Some fractional differential equations by Sumudu transform* International Journal of Scientific and Innovative Mathematical Research, (IJSIMR), 3(2), 615-619 (2015).
- [6] Bodkhe.D .S and Panchal. S. K., *On Sumudu transform of fractional derivatives and its applications to fractional differential equations*, Asian Journal of Mathematics and Computer Research, 11(1), 69 - 77 (2016).
- [7] Belgacem, F. B. M, Karaballi,A.A, and Kalla S.L. *Analytical investigations of the Sumudu transform and applications to integral production equations*, Mathematical Problems in Engineering, 3, 103-108 (2003).

- [8] Belgacem, F. B. M, Karaballi A.A. *Sumudu transform fundamental properties investigations and applications*, Journal of Applied Mathematics and Stochastic Analysis, Article ID 91083, 1-23 (2006).
- [9] Bulut, H. , Baskonus, H. M, Belgacem, F.B.M, *The Analytical solution of some fractional ordinary differential equations by the Sumudu transform method*, Abstract and Applied Analysis, 6pp (2013).
- [10] H.Bulut, H.Mellumet Baskonus, Seyma Tuluçe, *Homotopy Perturbation Sumudu Transform Method for Heat Equations*, Mathematics in Engineering, Science and Aerospace (MESA), 4(1), 49-60 (2013).
- [11] H.Bulut, H.Mellumet Baskonus, Seyma Tuluçe, *The Solutions of Homogeneous and Nonhomogeneous Linear Fractional Differential Equations by Variational Iteration Method*, Acta Universitatis, 36, 235-243 (2013).
- [12] H.Bulut, H.Mellumet Baskonus, Seyma Tuluçe, *The Solution of Wave Equations by Sumudu Transform Method*, Journal of Advanced Research in Applied Mathematics, 4(3), 66-72, (2012).
- [13] H.Bulut, H.Mellumet Baskonus, Seyma Tuluçe, *Homotopy Perturbation Sumudu Transform Method for One and Two Dimensional Homogeneous Heat Equations*, International Journal of Basic and Applied Sciences, IJBAS-IJENS, 12(1), 6-16 (2012).
- [14] Chaurasia, V. B. L, Dubey, R. S , Belgacem, F. B. M. *Fractional radial diffusion equation analytical solution via Hankel and Sumud transforms*, Mathematics In Engineering, Science And Aerospace, 3(2), 1-10 (2012).
- [15] Debnath, L. *Recent applications of fractional calculus to science and engineering*, International Journal of Mathematics and Mathematical Sciences, 54, 3413-3442 (2003).
- [16] Debnath, L and Bhatta. D, *Integral transforms and their applications*, Chapman and Hall /CRC, Taylor and Francis Group, New York (2007).
- [17] Seyma Tuluçe Demiray, Hasan Bulut and Fethi Bin Muhammad Balgacem, *Sumudu Transform Method for Analytical Solutions of Fractional Type Ordinary Differential Equations*, Mathematical Problems in Engineering, Article ID 131690, 6 pages, (2015).
- [18] Erdelyi(ed), A. *Higher Transcendental Function* , vol. 3. McGraw-Hill, New York, (1955).
- [19] Kilicman, A. , Eltayeb, H. and Agarwal, P.R. *On Sumudu transform and system of differential equations*, Abstract and Applied Analysis, Article ID 598702, 11 pp (2010).

- [20] Kilicman, A. and Eltayeb, H. "On the applications of Laplace and Sumudu transforms," Journal of the Franklin Institute, 347(5), 848-862 (2010).
- [21] Kilbas, A. A, Srivastava, H. M and Trujillo, J. J. *Theory and Application of Fractional Differential Equations* , Elsevier, Amsterdam, (2006).
- [22] Kataetbeh, Q. D and Belgacem, F. B. M. *Applications of the Sumudu transform to differential equations* , Nonlinear Studies, 18(1), 99-112 (2011).
- [23] Loonker, Deshna and Banerji, P. K., *Fractional integrals and derivatives for Sumudu transform on distribution spaces*, International Journal of Applications and Applied Mathematics, 7(1), 188-200 (2012).
- [24] Miller, K. S and Ross B. *An Introduction to the Fractional Calculus and Fractional Differential Equations* , John Wiley and Sons, (1993).
- [25] J.A.Nanware, G.A.Birajdar. *Methods of Solving Fractional Differential Equations of Order  $\alpha$  ( $0 < \alpha < 1$ )*, Bulletin of Marathwada Mathematical Society, 15(2), 40-53 (2014).
- [26] J.A.Nanware, D.B.Dhaigude, Monotone Technique for Finite System of Caputo Fractional Differential Equations with Periodic Boundary Conditions, Dyn. Conti., Disc. Impul. Sys., 22(1), 13-23 (2015)
- [27] Podlubny, I. *Fractional Differential Equations*, Academic Press, San Diego, (1999).
- [28] Weerakoon, S. *Application of Sumudu transform to partial differential equations*, International Journal of Mathematical Education in Science and Technology, 25(2), 277-283 (1994).
- [29] Weerakoon, S. *Complex Inversion Formula for Sumudu Transform* , International Journal of Mathematical Education, Science and Technology, 29(4), 618-621 (1998).
- [30] Watugala, G. K. *Sumudu Transform- an Integral transform to solve differential equations and control engineering problems*, International Journal of Mathematical Education in Science and Technology, 24(1) (1993).
- [31] Watugala, G. K. *Sumudu transform: a new integral transform to solve differential equations and control engineering problems*, Mathematical Engineering in Industry, 6(4), 319-329 (1998).