

Correlation Studies to Evaluate the Effect of Selected Soil Physical and Chemical Properties on Phosphate Availability in Hadejia-Nguru Floodplains, Nigeria

ABSTRACT

The purpose of the study was to evaluate the effect of selected soil physical and chemical properties on phosphate (P) availability in Hadejia-Nguru floodplains using correlation matrix. Three different floodplains were strategically selected. Disturbed soil samples were collected at three different sampling points on each floodplain and at three different soil depths 0-35, 35-70, 70-105 cm. Soil parameters were determined using standard laboratory procedures; Particle size distribution using hydrometer method, soil organic carbon using Walkley-Black Method, Electrical conductivity using conductivity meter, effective cation exchange capacity (ECEC) by summation method. Soil solution pH using glass electrode pH meter, available phosphorus using Bray 1 extraction method. The soil clay content range between 10 to 28 % clay. The soil organic carbon ranged between 1.5 to 7.7 gkg⁻¹. The soil EA values ranged between 0.9 to 1.3 cmol(+)kg⁻¹. The Ca²⁺ content range between 1.11 to 8.20 cmol(+)kg⁻¹. The exchangeable magnesium was between low to moderate, between 0.38 to 2.65 cmol(+)kg⁻¹. Values obtained in the floodplains were generally very low 0.010 to 0.026 cmol(+)kg⁻¹. The exchangeable sodium ranged between 0.87 to 1.24 cmol(+)kg⁻¹. There was a significant positive correlation between soil available P and OC content ($r = 0.542^ P < 0.05$), also between soil available P and pH ($r = 0.525^* P = .05$), supported by a significant negative correlation between E.A and available P ($r = - 0.604^* P = .05$). The soils manifest high significant negative correlation between soil available P and Ca²⁺ ($r = - 0.630^{**} P < 0.01$) as well as ECEC ($r = - 0.586^{**} P < 0.01$). Thus, Soil solution pH, exchangeable acidity (H⁺+Al³⁺), calcium ions Ca²⁺ and ECEC were the major soil properties that controls P availability in Hadejia-Nguru floodplains.*

Key Words: Correlation, Soil Properties, Sorption, Floodplains, Phosphate Availability.

1.1 INTRODUCTION

Soil physical and chemical properties such as; clay content, organic matter (O.M), cation exchange capacity (C.E.C) and soil pH affect phosphate availability greatly as discussed by Hamoud, [11]. Soils with high clay content have a greater capacity to loosely adsorb and retain P making it potentially available for plant uptake as discussed by Yan, [26], Hamoud, [11], Chen, and Arai, [5]. High clay content and/or Organic matter (OM) increases soil Cation Exchange Capacity (CEC) as discussed by Hamoud,[11]. and higher CEC contributed from O.M can enhance P availability greatly as discussed by Antonangelo, [4], Hamoud,[11], due to its chelating ability Orji, [20]. The naturally synthesized hydroxyl and carboxyl ligands in the humic, fulvic and low-molecular weight organic acids affect phosphate availability by being attracted to some of the reactive sites on the soil matrix surfaces thereby reducing strong P fixation as discussed by Ahmed [2] and Li, [15]. The concentrations of H_2PO_4^- and HPO_4^{2-} in the soil largely depends on soil solution pH as discussed by Johan, [12]. The objectives of this research were to evaluate the impact of selected parameters controlling phosphorus availability in Hadejia-Nguru Floodplains. The major problem associated to flooded soils is formation of acidity due to aluminum hydrolysis and transformation of iron (III) in to Iron (II) which reacts with phosphate ions and make them unavailable for plants uptake, where this problem exists organic matter application can help solve or reduce the problem by complexing and blocking the reactive sites of Al^{+3} and Fe^{+2} ions thereby inhibiting strong fixation of P ions on the aluminum and iron ions and make the P ions readily free in solution. Most of the recent researches as discussed by Goni, [10] and Mohammed, [17]) conducted in Hdejia-Nguru floodplains focus more on fertility status, pH, exchangeable bases and CEC of the wetlands. Hence, the need for this research to close the gap.

2.1 MATERIALS AND METHODS

2.1.1 Site Description

The research was conducted in Hadejia-Nguru floodplains in Sudan savanna of Nigerian. Three different floodplains namely northern floodplains located at 12.450°N and 10.044°E, western floodplains 12.405°N and 10.100°E and southern floodplains 12.447°N and 10.130°E of the Hadejia-Nguru river basin were selected. Disturbed soil samples were collected using W-shaped sampling strategy at three different sampling points and three different soil depths 0-35, 35-70, 70-105 cm from each of the floodplains, then composited to a total of 9 samples. The soil samples were air dried in the laboratory, crushed, passed through 2mm mesh sieve, labeled and stored in dark polyethene leathers for further laboratory analysis. Statistical package Software version 9.3. SAS, [23] was use for the data analysis

2.1.2 Extraction and Analytical Procedures

The following procedures were used for the soil physical and chemical analysis; Particle size distribution using hydrometer method described by Sedlackova, [21], soil organic carbon using Walkley-Black method described by Mustapha, [18], Electrical conductivity using conductivity meter method described by Mizuhata, [16], Exchangeable bases were extracted using 1M ammonium acetate extraction method, exchangeable bases (K^+ and Na^+) were determined using flame photometer, (Mg^{++} and Ca^{++}) were determined using A.A.S described by FAO, [7]. Exchangeable acidity (H^+ and Al^{+++}) were determined by titration method and finally ECEC was estimated by summing up exchangeable bases and exchangeable acidity described by FAO, [7]. Soil solution pH was determined using glass electrode pH meter described by FAO, [9] Soil available phosphorus was extracted using Bray No.1 described by FAO, [7] and the P in solution was analyzed using spectrophotometric technique described by FAO, [7].

3.1 RESULTS AND DISCUSSION

3.1.1 Soil Clay Content

The statistical data obtained revealed that, the soil clay content (Table 1.0 and 2.0) recorded in the floodplains were within the range of 10 to 28 % clay. Soils with the highest clay contents were recorded in the surface soils of the southern and northern floodplains (Table 1.0 and 2.0) having 26 and 28 % clay. Consistent increase down the soil profile was observed in all the floodplains except in northern floodplain.

Table 1: The Particle-Size Distribution of Hadejia-Nguru Floodplains

Sampling Locations	Depth (cm)	% Clay	% Silt	% Sand	Textural Class
NFpHJRB	0-35	14.00	51.00	35.00	Silt loam
	35-70	14.00	35.00	52.00	Loam
	70-105	12.00	38.00	50.00	Loam
WFpHJRB	0-35	26.00	31.00	43.00	Loam
	35-70	16.00	32.00	52.00	Loam
	70-105	13.00	25.00	62.00	Loamy sand
SFpHJRB	0-35	28.00	28.00	44.00	Clay loam
	35-70	20.00	22.00	58.00	Sandy loam
	70-105	10.00	39.00	51.00	Loam

N = northern, W = western, S = southern, Fp = Floodplain, HJRB = Hadejia-Jama'are river basin

Table 2: Selected Soil Physical and Chemical Properties of Hadejia-Nguru Floodplains

Loc.	Depth(cm)	%Clay	O.C	pH	CaCl ₂	EA	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	ECEC	EC(dSm ⁻¹)	Av.P
		(gkg ⁻¹)				(cmol(+)k g ⁻¹)					(mgkg ⁻¹)		
NF	0-35	14	5.2	5.0	0.9	0.35	0.34	0.017	1.02	2.68	1.02	19.6	
	35-70	14	4.1	5.1	0.9	0.77	0.40	0.017	1.03	3.17	1.21	14.2	
	70-105	12	2.9	4.8	1.0	0.51	0.31	0.010	0.97	2.79	0.57	12.4	
WF	0-35	26	7.7	5.1	1.3	0.35	0.35	0.011	1.24	3.45	1.02	16.2	
	35-70	16	4.6	5.1	1.1	0.71	0.43	0.026	0.90	3.26	0.92	14.6	
	70-105	13	2.9	4.4	1.3	0.55	0.22	0.019	1.14	3.00	0.89	10.5	
SF	0-35	28	2.5	4.1	1.3	0.35	0.32	0.012	1.10	3.09	0.99	13.4	
	35-70	20	1.9	5.3	0.9	0.66	0.41	0.017	0.90	3.04	0.86	9.4	
	70-105	10	1.5	4.1	1.1	0.60	0.33	0.018	0.87	2.08	0.87	8.0	

Loc.= Location, **NF** = Northern Floodplains, **WF** = Western Flood Plains, **SF** = Southern Floodplains,

SiL = Silt Loam, **SL** = Sandy Loam, **LS** = Loamy Sand, **L** = Loamy Soil, **CL** = Clay Loam

3.1.2 Soil Organic Carbon

The soil organic carbon of the floodplains was within the range of 1.5 to 7.7 gkg⁻¹. All the floodplains reveal drastic decrease down the profile. Surface soils of northern and western floodplains have higher organic carbon content (5.2 and 7.7) compared to the southern flood plain with (2.5) and similar trend was observed in all the three soil depths with southern flood plain having very low organic carbon content.

3.1.3 Soil Solution pH

The pH of the floodplains (Table 2.0) varied between strongly acidic (pH 4.1) to moderately acidic (pH 5.5). Southern flood plain has low pH 4.1 compared to northern and western floodplains (pH 5.0 and 5.1). Mohammed, [17] reported a pH of 5.7 in Hadejia-Jama'are soils which corroborate with some of the values recorded in this work.

3.3.4 Exchangeable Acidity (EA)

The floodplains appeared to have a range of EA values between 0.9 to 1.3 cmol₍₊₎kg⁻¹. All the floodplains manifest increase in E.A down the soil profile except southern floodplain that manifest an inconsistent pattern. Shehu, [22] reported values < 1.0 cmol₍₊₎kg⁻¹ in Nigeria Sudan savanna which agree with the values recorded in this work.

3.1.5 The Exchangeable Calcium (Ca²⁺)

The Ca²⁺ content of the floodplains were within the range of 1.11 to 8.20 cmol₍₊₎kg⁻¹ which is generally low. Abdu and Udofot, [1] show that, Ca is a dominant cation in the exchange site of savanna drylands. Goni, [10] reported 1.240 meq/L in the same wetlands of Hadejia-Jama'are. Shehu, [22] reported 2.0 to 5.0 cmol₍₊₎kg⁻¹ in dryland soils. Mohammed, [17] reported a range of values of 3.44 to 4.31 cmol₍₊₎kg⁻¹ in Hadejia- Nguru wetlands, Nigeria.

3.1.6 The Exchangeable Magnesium (Mg^{2+})

The exchangeable magnesium concentration recorded in the floodplains ranged between 0.38 to 2.65 $cmol_{(+)}kg^{-1}$. Values obtained in all the locations were between low to moderate, since values $\leq 0.4 cmol_{(+)}kg^{-1}$ is considered low (Leo, [14], Daniel, [6]. Mohammed, [17] reported a range of values of 1.752 to 2.158 $cmol_{(+)}kg^{-1}$ in Hadejia-Nguru wetlands, Nigeria, their report agree with the values recorded in this research work. Shehu, [22] reported a range of 0.3 to $> 1.0 cmol_{+}kg^{-1}$. some of the values he reported agree with the values recorded in this research work. Goni, [10] reported 0.358 meq/L in the same wetlands of Hadejia-Jama'are.

3.1.7 The Exchangeable Potassium (K^{+})

Values obtained in the floodplains were generally very low 0.010 to 0.026 $cmol_{(+)}kg^{-1}$ attributed to non-exchangeable behavior of K^{+} , often trapped in the inter-layer of clay minerals, such that very little amount is freely available. Goni, [10] reported values (1.1 meq/L) which were higher compared to the values obtained in this work. Mohammed, [17] reported a range of values of 0.356-0.692 $cmol_{(+)}kg^{-1}$ in Hadejia-Nguru wetlands, Nigeria, values reported in their work are higher than the values recorded in this work, the differences may be due to spacial variability of the soils even within the same environment as well as the sampling period.

3.1.8 The Exchangeable Sodium (Na^{+})

The exchangeable sodium concentration of the floodplains (Table 2.0) ranged from 0.87 to 1.24 $cmol_{(+)}kg^{-1}$. The soils manifest no evidence of sodicity, because for a soil to be considered sodic it must have exchangeable sodium percentage (ESP) $>15 \%$, $EC < 4 (dSm^{-1})$, $pH \geq 8.2$ and/or manifest visible effects on the soil structure FAO, [8]. Shehu,[22] reported

very low values (0.1 to 0.3 $\text{cmol}_{(+)}\text{kg}^{-1}$) in Nigerian Sudan Savanna soils. Yakubu, [25] reported 0.01 to 0.12 $\text{cmol}_{(+)}\text{kg}^{-1}$ in Nigerian Guinea Savanna. Mohammed, [17] reported a range of values of 0.021 to 0.033 $\text{cmol}_{(+)}\text{kg}^{-1}$ in Hadejia-Nguru wetlands, these values are extremely low compared to the values reported in this research work.

3.1.9 Correlation of the Selected Soil Physical and Chemical Properties with Available P

The results obtained (Table 3.0) contains both positively and negatively significant correlation values between phosphate availability, soil physical properties and soil chemical properties. A significant positive correlation was observed between soil available P and soil solution pH ($r = 0.525^* P = .05$) suggesting that soil available P will become readily available as pH improve from strongly acidic towards the neutral region and as the pH reversed to strongly acidic region available P will become more strongly tied-up and un-available for plants uptake. A significant positive correlation was observed between soil available P and OC content ($r = 0.542^* P = .05$), this also shows that increase in OC content will leads to corresponding increase in P availability in the soil and vice- versa. Some researchers Tsozue, [24], Amenkhienan, and Isitekhale,[3], Kadiri, [13] reported significant positive correlations between available P and organic carbon in a Typic Dystrandept soils of Cameroon western highland ($r = 0.87, P = .05$), and on cretaceous sediments, shale and quaternary alluvium in Edo state, Nigeria ($r = 0.980^{**} P < 0.01$) and also in different land use type of two agro-ecological zones of Nigeria respectively ($r = 0.53^{**} P < 0.01$). A significant negative correlation was observed between available P and exchangeable acidity (H+Al) ($r = - 0.604^* P = .05$). A high significant negative correlation was observed between available P and Ca^{2+} ($r = - 0.630^{**} P < 0.01$). Reports by (Amenkhienan, and Isitekhale, [3], Ojorbor, and Aimufia, [19] reveal a high positive correlation between available P and Ca^{2+} ions in cretaceous sediments, shale and quaternary alluvium in Edo state, Nigeria ($r = 0.969^{**} P <$

Table 3: Correlation Matrix of the Selected Soil Physical and Chemical Properties on Available Phosphorus

	Clay	pH CaCl ₂	EC	Av.P	Na ⁺	K ⁺	Ca ²⁺	Mg ²⁺	H+Al	ECEC	OC
Clay											
pH	- 0.635** 0.0034										
EC	0.190	0.101									
Av.P	- 0.390	0.525* 0.0392	- 0.412								
Na ⁺	0.092	0.150	0.511* 0.0302	0.010							
K ⁺	- 0.029	- 0.189	- 0.401	- 0.048	0.009						
Ca ²⁺	0.685*** 0.0017	- 0.682*** 0.0018	0.559* 0.059	- 0.630** 0.0050	0.095	0.1803					
Mg ²⁺	0.450	0.313	0.006	0.204	0.188	- 0.100	- 0.177				
H+Al	0.582** 0.0112	0.910** 0.016	0.208	- 0.604* 0.0479	0.062	0.167	- 0.034	- 0.083			
ECEC	0.696*** 0.0013	- 0.624** 0.0056	0.119	- 0.586** 0.0106	0.294	0.348	0.970*** < 0.0001	- 0.081	0.098*** < 0.001		
OC	0.571* 0.0013	0.442	0.370	0.442	0.055	0.430	- 0.408	- 0.209	- 0.245	0.431	

EC = electrical conductivity, Av. p = available phosphorus, OC = organic carbon, ECEC = effective cation exchange capacity,

H+Al = hydrogen plus aluminum ions, pH = hydrogen ions potential of the soil solution. EC = electrical conductivity

0.01) and in Fadama soils of Asaba, Delta state, Nigeria, ($r = 0.0665^* P < 0.01$). A high significant negative correlation was observed between available P and ECEC ($r = - 0.586^{**} P < 0.01$) in Hadeja-Jama'are floodplains and in contrary, a very strong positive correlation was reported by Amenkhienan, and Isitekhale, [3] $r = 0.944^{**} P < 0.01$ on cretaceous sediments, shale and quaternary alluvium in Edo state, Nigeria. Soil clay content shows a very significant positive correlation with pH CaCl₂ ($r = 0.635^{**} P < 0.01$), EA ($r = 0.582^{**} P < 0.01$) and a very high significant positive correlation with ECEC ($r = 0.696^{***} P < 0.001$) and with Ca²⁺ ($r = 0.685^{***}$). Electrical conductivity (EC) correlates significantly positive with Na⁺ ($r = 0.511$) and Ca²⁺ ($r = 0.259^*$) while soil solution pH correlate negatively and very significantly negative with Ca²⁺ ($r = - 0.682^{***} P < 0.001$) but EA reveal very significantly negative correlation with pH, indicating that an increase in EA will consequently leads to corresponding decrease in soil solution pH and make the soil solution more acidic. A high significant positive correlation was observed between ECEC and Ca²⁺ content of the study area ($r = 0.970^{***} P < 0.001$), suggesting that an increase or decrease in Ca²⁺ content of Hadeja-Jama'are floodplains is directly proportional to ECEC.

4.1 CONCLUSION

Base on the data presented in this study, the following conclusions were made: Soil solution pH, exchangeable acidity (H⁺+Al³⁺), calcium ions Ca²⁺ and estimated cation exchangeable capacity ECEC were the major soil properties that controls P availability in Hadejia-Nguru floodplains as at the time this research was conducted, because the correlation matrix revealed a significant positive correlation between soil available P and pH ($r = 0.525^* P = .05$), suggesting that soil P becomes more available as pH improve from strongly acidic towards the neutral region vice-versa. This was also supported by the significant negative correlation ($r = - 0.604^* P = .05$) between E.A

and available P, indicating that as E.A increases P availability decreases. The soils manifest high significant negative correlation between soil available P and Ca^{2+} ions as well as ECEC ($r = -0.586^{**}$ $P < 0.01$) indicating that, high amount of Ca^{2+} ions increase soil ECEC and an increase in ECEC will lead to increase in high P sorption. The soils of the Hadejia-Nguru floodplains were strongly to moderately acidic as at the time this research was conducted. The soils exhibit low clay content. The soils have very low exchangeable K^+ associated to non-exchangeable behavior of K^+ often trapped in the inter-layer of clay minerals.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

8.1 REFERENCES

1. Abdu N. and Udofot E. A. (2015) Fifteen-year Fallow Altered the Dynamics of Soil Phosphorus and Cationic Balance in a Savanna Alfisol. *Archives of Agronomy and Soil Science*. 61, 645-656
2. Ahmed, A.A., Leinweber, P., and Kuhn, O. (2023). Advances in Understanding the Phosphate Binding to Soil Constituents: A Computation Chemistry Perspective. *Science of the Total Environment*, 887, 163692
3. Amenkhienan, B.E. and Isitekhale, H.H.E (2021). Physicochemical Properties of Selected Soils Formed on Cretaceous Sediments, Shale and Quaternary Alluvium in Edo State, Nigeria. *Journal of Current Opinion in Crop Science* 2(2), 192-201
4. Antonangelo, J.A., Culman, S., and Zhang, H. (2024) Comparative Analysis and Prediction on Cation Exchange Capacity Via Summation: Influence of Biochar Type and Nutrient Ratios. *Frontiers*. 4, 2024.

5. Chen, A. and Arai, Y. (2024) A Review of the Reactivity of Phosphate Controlled by Clays and Clay Minerals: Implication for Understanding Phosphate Mineralization in Soils. *Clays and Clay Minerals*. 17(2)
6. Daniel C.J., Frid, L., Sleeter, B.M and Fortim, M.J. (2016) State and Transition Simulation Model: A Framework for Forecasting Landscape Change. *Methods in Ecology and Evaluation* (7),11
7. Food and Agriculture Organization United Nations F.A.O (2002). *Procedure for Soil Analysis*. Van Reeuwijk, L.P (editor) 6th Ed. Wageningen. International Soil Reference and Information Centre. Wageningen. The Netherlands Pp. 9-14
8. Food and Agriculture Organization United Nations F.A.O (2019). Sodic Soils and their Management. Page/Site visited on 7/6/2019 Retrieved from www.fao.org/3/x5871e/5871e05...
9. Food and Agriculture Organization United Nations F.A.O (2021). *Standard Operating Procedure for Soil pH determination*. Rome
10. Goni, I., Vassolo. S., Sherif, M., Aji, M., Bura, B.B. and Ibrahim, Y. (2023). Geochemical and isotopic studies in parts of the Hadejia-Jamaare- Komadugu-Yobe Basin, northeast Nigeria. *Hydrogeology Journal*. 32, 113–129
11. Hamoud, Y.A., Shaghaleh, H., Zhang, K., Okla, K.M., Alaraidh, I.A., and S.M., Elgawad, H.A. (2024). Increasing Soil Clay Content Increases Soil Phosphorus Availability and Improves the Growth, Physiology, and Phosphorus Uptake of Rice Under Alternative Wetting and Mild Drying Irrigation. *Environmental Technology & Innovation*. 35
12. Johan, P.D., Ahmed, O.H., Omar, L. and Hasbullah, N.A. (2021) Phosphorus Transformation in Soils Following Co-Application of Charcoal and Wood Ash. *Agronomy*. 11(10),2010. <https://doi.org/10.3390/agronomy11102010>

13. Kadiri, W.O.J., Ogunleye, K.S., Fasina, A.S. and Babalola, T.S. (2022) Soil properties affecting soil organic carbon stock of different land use type of two agro-ecological zones of Nigeria. *Journal of Tropical Agriculture, Food, Environment and Extension*. 21,37-48
14. Leo, E., Nathan, S., and Morteza, M., (n.d) *Understanding Numbers on your Soil Test Report*. University of Arkansas, USA. Retrieved from <https://www.uaex.edu>>PDF Site visited 29/6/2019
15. Li, C., He, M., Xin, C. and Zhang, Z. (2024). Phosphate Desorption Regulates Phosphate Fraction Dynamics In Aggregates of Revegetated Ecosystem. *Journal of Environmental Management*. 361, 121238
16. Mizuhata, M. (2022). Electrical Conductivity Measurement of Electrolyte Solution. Review. *Electrochemistry*. 90(10), 102011
17. Mohammed, S.E., Aderibigbe, D.O., Saeed, M.D., Nelana, S.M., Musa, A. and Ayanda, O.S. (2024). Seasonal Quality and Statistical Assessment of Physicochemical Properties and Base Saturation of Hadejia-Nguru Wetland Soils, Northwest/Eastern Nigeria. *Water practice and Technology*. 19(3), 761-774.
18. Mustapha A.A (2020). Comparison of Different Method for the Assessment of Soil Organic Carbon in Northern Guinea Savanna Zone of Nigeria. *Sustainable Agriculture and environment*. 18(1), 36-41
19. Ojorbor, S.A., and Aimufia, P.N (2022). Fertility Status of Fadama soils of Asaba, Delta state, Nigeria. *Journal of Soil Science*. 32(1), 72-76
20. Orji, J. E., Igwe, C. A. and Ezeaku, P. I. (2023). *Effect of Climate Change on Soil Organic Carbon Storage in Four Land Use Types in Abakaliki, South Eastern Nigeria*. Johnbosco, C., Egbueri, J.C., Ighalo, J.O. and Pande, C.A.B. (Ed). *Climate Change Impacts on Nigeria Environment and Sustainable Development*. Pp. 135

21. Sedlackkova, K., Sevelova, L., Igaz, D. and Aydin, E. (2024). Determination of Particle Size Distribution: Comparison of Standard Hydrometer Method and Laser Diffraction Analysis for Use in Forestry. *Forest*. 15(2), 327
22. Shehu, B.M., Jibrin, J.M., and Samndi, A.M. (2015) Fertility Status of Selected Soils in the Sudan Savanna Biome of Northern Nigeria *International Journal of Soil Science* 10(2), 74-84
23. Statistical Analysis System SAS (2011). Statistical package Software version 9.3.SAS Institute, Cary, North Carolina.
24. Tsozue D., Tematio, P., Azinwi, T.P (2016) Relationship Between Soil Characteristics and Fertility Implication in two Typical Dystrendept Soil of the Cameroon Western High Land. *International Journal of Soil Science*. 3(2), 36-40
25. Yakubu, A. (2016). *Interaction of Microbial, Organic and Inorganic Sources of Phosphorus on the Performance of Maize (Zea mays L.) in Two Soils of Different Fertility*. (Project for Bachelor's Degree in Agriculture, Ahmadu Bello University)
26. Yan, X., Yang, W., Chen, X., Wang, M., Wang, W., Ye, D. and Wu L. (2020) Soil Phosphorus Pools, Bioavailability and Environmental Risk in Response to the Phosphorus Supply in the Red Soil of Southern China. *Int J Environ Res Public Health*. 17(20), 7384.