

# Unraveling Genetic Variability, Correlation and Path Analysis for Yield and Its Components in Barley(*Hordeum vulgare* L.)

## Abstract

The current study aimed to assess the genetic variability of yield and yield-related traits while examining the direct and indirect effects of trait interactions. Conducted at a research farm in Chaudhary Charan Singh University, Meerut, Uttar Pradesh, India, the experiment involved fourteen parents and their BC<sub>1</sub>F<sub>1</sub> crosses, analyzed using a randomized block design (RBD). Significant differences were observed among the varieties for most traits, indicating a broad range of mean values and diversity. Both phenotypic and genotypic coefficients of variance (PCV and GCV) were generally low, with biological yield showing the highest values (26.57 for GCV and 40.48 for PCV). PCV values were slightly higher than GCV values. Traits such as seed per spike, biological yield, flag leaf breadth, days of heading, and days of maturity exhibited relatively high heritability. Notably, grain yield showed high heritability and genetic progress, making it a favourable selection indicator. Several traits, including days of anthesis, days of maturity, and spike length, demonstrated significant positive correlations with yield while also showing significant negative correlations with grain filling period and biological yield. Conversely, these traits showed highly non-significant positive correlations with harvest index, plant height, thousand seeds weight, and chlorophyll content and highly non-significant negative correlations with tiller number and seeds per spike.

**Keywords:** Genotypic and Phenotypic variance, Heritability, Coefficient, Variability.

## 1. Introduction

Barley belongs to the genus *Hordeum* and Poaceae family (Von Bothmer *et al.*, 1995; Kling and Hayes, 2004)<sup>[19,10]</sup>. During the second half of the second millennium, barley arrived in China. One of the world's oldest food crops is barley (*Hordeum vulgare* L.). Since the early stages of agricultural developments 8,000-10,000 years ago, it has been a significant cereal crop (Giles and Bothmer, 1985 & Bourne, 1987)<sup>[3,4]</sup>. It is a commercially significant cereal crop, ranking fourth in the world after wheat, rice, and maize in quantity produced and cultivated area (FAO, 2014). Barley is native to the Eastern Mediterranean, where plants are subjected to various abiotic stresses in the field. It is grown in many places where the climate could be more favourable. Though it has a lower commercial value than wheat, it replaces wheat in dry areas where water is scarce. It is known as the poor man's crop because of its minimal input

requirements and superior tolerance to rainfed conditions (Verma *et al.*, 2011)<sup>[18]</sup>. Barley production in the world totals 292.9 million tonnes, with Europe producing the most (59.6%), followed by Asia (14.9%). The Russian Federation is the leading producer, with a total output of around 20.02 million tonnes, whereas India is ranked fourteenth (USDA, 2015). In 2017, India's barley production was 1.75 million tonnes, but according to 2008-09 statistics, barley is planted on 0.71 million hectares with a production of 1.69 million tonnes and a yield of 2394 kg ha<sup>-1</sup>. Rajasthan has the most area (0.29 million ha) and production (0.89 million t) of barley, whereas Haryana has the highest yield (3491 kg ha<sup>-1</sup>). Cultivated barley is a species of *Hordeum* that evolved from wild barley (*Hordeum spontaneum*), which can still be found in the Middle East. Cultivated and wild barley have fourteen chromosomes (2n=14) and are diploid species. *Hordeum vulgare* L (Jain, 1960)<sup>[8]</sup>. It is the only cultivated species with two phenotypic variants, six-rowed (*Hordeum vulgare*, *H. hexastichum*) and two-rowed (*Hordeum vulgare*, *H. hexastichum*) (*H. distichum*). They have the same chromosomal number (2n=14) and may intercross freely to create fertile hybrids despite their spike morphological variances (Poehlman, 1987 & Singh, 1985)<sup>[5,13]</sup>. Barley contains a great deal of genetic variety used to classify the species. There are many different methods to categorize barley. Identifying whether the spike has two, four, or six rows of spikelet's is one technique to classify barley. Most cultivated barley has six rows, while wild barley has two.

## **2. Materials and methods**

### **2.1. Experimental site, Data collection, material and procedures**

The experimental material for the present study was obtained from *Eternal University, Baru Sahib, Dist. Sirmour, H.P.* The present investigation was conducted at the Research Farm and Molecular Laboratory of *Dept. of Genetics and Plant Breeding C.C.S. University Campus, Meerut (UP) during the year 2020-2021*, with three replications; the trial was set up in a Randomized Block Design (RBD). The Standard Evaluation System for barley developed by the Indian Institute of Wheat and Barley Research was used to make observations and data records for all attributes investigated. Five sample plants were randomly selected from each plot in the middle three rows. Observations were recorded on seventeen quantitative traits Days to heading (DTH), Date of anthesis (DTA), Date of maturity (DTM), Grain filling period (GFD), Average plant height (APH), Flag leaf length (FL), Flag leaf width (FW), spike length (SL), Average tiller number, (ATN), No. of grains per spike (NG), biological yield (BY), Germination percentage (GER), Chlorophyll content (CC), grain yield (GY), thousand-grain yield (TGW), Harvest index

(HI), were used for assessing the genetic diversity and characters association among these genotypes of barley.

Table 1: A list of barley cultivar used for evaluation and crossing in the present study

S.No.	Nameofcultivar/varieties	2rows/6rows	Remarks
1.	IITR-39	6rows	Hullessbarley
2.	PL-830	6rows	Hulledbarley
3.	PL-172	6rows	Hulledbarley
4.	PL-707	6rows	Hulledbarley
5.	PL-419	6rows	Hulledbarley
6.	PL-751	6rows	Hulledbarley
7.	PL-426	6rows	Hulledbarley
8.	PL-758	6rows	Hulledbarley
9.	IITR-104	6rows	Hulledbarley
10.	IITR-38	2rows	Hulledbarleyanda nthocyaninrich
11.	IITR-35	2rows	Hulledbarley
12.	VIJY-102	2rows	Hulledbarley
13.	DWRUB-52	2rows	Hulledbarley
14.	PL-838	2rows	Hulledbarley

## 2.2.Statistical analysis

Analysis of variance was performed using the plant breeding statistical program SPSS software. The genotypic and phenotypic variance, genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability in broad sense ( $h^2_b$ ), genetic advance in percentage of mean (GA), genotypic correlation coefficients ( $r_g$ ) and phenotypic correlation coefficients ( $r_p$ ), genotyping and phenotyping path analysis were estimated following (Singh & Chaudhary, 1985)<sup>[15]</sup>. The estimates of the Genetic Coefficient of Variation (GCV) and Phenotypic Coefficient of Variation (PCV) were categorized into low, medium, and high according to the classification proposed by Sivasubramanian and Madhavamenon in 1973<sup>[16]</sup>. Heritability in a broad sense and genetic advance were calculated according to methods given by (Allard, 1960; Singh and Chaudhary, 1985)<sup>[2,15]</sup>. Path coefficient analysis was done using R software.

## 3. Results and discussion

Creating genetic variability and selecting key traits are crucial activities every plant breeder should undertake to enhance yield and other desirable agronomic characteristics. However, to effectively select, it is essential to have information on the available genetic

variation among barley genotypes. Thus, effective selection depends not only on estimating genetic variation among genotypes but also on the proportion of heritable variation and the expected genetic gain that would be obtained (Falconer & Mackay, 1996; Singh, 2000)<sup>[6,15]</sup>. Heritable variation is helpful for permanent genetic improvement (Singh, 2000)<sup>[15]</sup>. Heritability broadly estimates the ratio of total genetic variance, including additive, dominant and epistatic variances to the phenotypic variance (Falconer and Mackay, 1996; Riaz and Chaudhary, 2004)<sup>[6,14]</sup>.

The ANOVA indicated significant differences among the cultivars for Days to heading (DTH), Date of anthesis (DTA), Date of maturity (DTM), Grain filling period (GFD), Average tiller number, (ATN), No. of grains per spike (NG), biological yield (BY), Germination percentage (GER), grain yield (GY), thousand-grain yield (TGW). The analysis of variance also revealed highly significant differences among the test genotypes for all the traits studied. The mean sum sequences due to test genotypes were highly significant for Date of maturity (DTM), Grain filling period (GFD), Average tiller number (ATN), No. of grains per spike (NG), biological yield (BY), Germination percentage (GER), grain yield (GY), thousand-grain yield (TGW).

The estimating of range (Minimum and maximum), Mean Standard Error, Critical difference (5%), Critical variance (1%), environmental variance, Genotypic variance, Phenotypic variance, Genotypic coefficient of variance, Phenotypic coefficient of variance, Heritability (Broad sense), Advancement in genetics as a percentage of the mean of 17 traits. The range of date of heading among the genotypes differs from 89 to 101 with a value of mean is 93.38 and CV is 3.51 per cent, days of anthesis among the genotypes differs from 93 to 105 as the range with a value of mean is 99.18 and CV is 4.14%, days of maturity among genotypes differ from 114 to 126 as the range with a value of mean is 118.87, CV is 3.09 followed by the other quantitative and qualitative traits (Table 2).

The highest GCV (26.57) and PCV (40.48) estimates for grain yield should have a high degree of genetic variation for this trait. However, a moderate level of differences among GCV (26.57) and PCV (40.48) for grain yield indicates the role of environmental variation for GCV and PCV estimates, thus having sufficient genetic variability for the improvement of these traits (Karzai, 2004)<sup>[10]</sup>. Some traits like the number of plants germinated, seeds per spike, thousand seeds weight, and average flag leaf length also had moderate values for GCV and PCV estimates, thus having sufficient genetic variability to improve these traits (Karthikeyan, 2010)<sup>[11]</sup>. A

meagre value of GCV and PCV for days of anthesis, days of heading and days of maturity harvest index, average spike length, chlorophyll content, leaf angle, days of anthesis, days of heading and days of maturity (Table.2) showed the tiny scope of improvement in the genotypes for these traits and similar results also reported by Yadav *et al.*, (2015)<sup>[21]</sup>.

Heritable variation is helpful for permanent genetic improvement (*Singh, 2000*)[15]. Heritability broadly estimates the ratio of total genetic variance, including additive, dominant and epistatic variances, to the phenotypic variance (Falconer & Mackay, 1996; Riaz and Chaudhary, 2004)<sup>[6,7,14]</sup>. The Highest heritability, along with the highest genetic advance, was recorded for biological yield, seeds per spike, thousand seeds weight and average plant height, similar results reported by (*Addisu et al.*, T., 2015)<sup>[11]</sup> (Table 2 & Figure 1). Hence these traits are under control for additive genes. These can be improved by selection based on phenotypic performance.

**Table2: Estimation of Mean, Range (Minimum & Maximum), Standard Error of mean, Critical variance (5%), Critical difference (1%), Environmental variance, Genotypic variance, Phenotypic variance, Environmental coefficient of variance, Genotypic coefficient of variance, Phenotypic coefficient of variance, Heritability (Broad sense), Genetic advance, Genetic advance as percentage of mean of seventeen different traits.**

Traits	Range		SEm	GV	PV	GCV	PCV	Heri.(BS)	G.advance	GA as % of mean
	Maxi.	Mini.								
DTH	101	89	1.25	2.46	7.13	1.63	2.86	0.34	1.9	2.03
DTA	105	93	1.47	1.7	8.2	1.32	2.89	0.21	1.22	1.23
DTM	126	114	1.1	1.67	5.29	1.09	1.93	0.32	1.5	1.26
GFD	26	9	1.51	2.79	9.63	8.48	15.76	0.29	1.85	9.41
ATN	9	2	0.58	0.23	1.24	12.09	28.33	0.18	0.42	10.62
AFL	23.63	8	1.39	0.56	6.35	5.52	18.58	0.09	0.05	3.37
AFW	1.73	0.4	0.12	0.04	0.08	18.24	26.93	0.46	0.27	25.45
APH	75.7	35.33	3.42	19.1	54.21	8.47	14.27	0.35	5.34	10.36
LA	2	1	0.27	0.01	0.22	5.77	34.4	0.03	0.03	1.97
BYL	240	45	19.27	843.26	1956.7	26.57	40.48	0.43	39.27	35.93
HI	41.18	16.12	2.48	2.91	21.31	6.29	17.03	0.14	1.3	4.79
ASL	11	4.67	0.56	0.21	1.14	6.3	14.75	0.18	0.4	5.54
SPS	74.66	14	5.92	84.42	189.58	19.88	29.79	0.45	12.63	27.33
NPG	40	3	5.69	-10.44	86.69	12.92	37.23	-0.12	-2.31	-9.24
CC	65.1	32.9	3.23	3.62	34.9	3.96	12.31	0.1	1.26	2.63
TW	72	22.1	2.83	81.74	105.72	18.58	21.14	0.77	16.38	33.66
YL	75	25	5.63	99.61	194.85	25.59	35.79	0.51	14.7	37.69





Figure 1: Estimation of genetic variability (GCV, PCV and Heritability)






Figure 2: Violin plot (A box or marker indicating the inter quartile range; and possibly all sample points for 16 different traits).

### 3.1. Estimations of genotypic and phenotypic correlation

The study identified significant correlations among various growth and yield-related traits in the crop. Days to 50% heading were positively correlated with days of anthesis and spike length but showed a non-significant positive correlation with days of maturity. Conversely, they were negatively correlated with grain filling period, flag leaf length, seeds per spike, chlorophyll content, and biological yield. Day 50 anthesis was positively correlated with spike length and non-significantly correlated with maturity, tiller number, plant height, harvest index, and thousand seed weight but negatively correlated with grain filling period, flag leaf length, seeds per spike, and chlorophyll content. Additionally, traits such as days of maturity, spike length, and days to 75% maturity were positively correlated with days of anthesis and spike

length and non-significantly correlated with days to maturity. Plant height, tiller number, and biological yield also displayed significant positive correlations with certain traits but showed negative correlations with others. These findings highlight complex trait interactions, which can inform breeding strategies for improving crop yield and quality.


UNDER PEER REVIEW

**Table3: Genotypic correlation and phenotypic correlation**

Traits	DTH	DTA	DTM	GFD	ATN	AFL	AFW	APH	BYL	HI	ASL	SPS	CC	TW	YL
<b>DTH</b>	1**	0.96**	0.10N	-0.67**	0.18NS	-0.69**	-0.19NS	0.23NS	-0.02NS	0.08NS	0.46**	-0.36**	-0.49 **	0.13 NS	-0.06NS
<b>DTA</b>	0.89 **	1 **	0.17 NS	-0.64 **	0.23 NS	-0.26 NS	0.02 NS	0.11 NS	0.04 NS	0.14NS	0.65 **	-0.56 **	-0.37 *	0.20 NS	0.04 NS
<b>DTM</b>	0.27 **	0.29 **	1 **	0.63 **	-0.07 NS	0.14 NS	0.04 NS	0.80 **	0.60 **	-0.4 *	0.33 *	0.21 NS	0.12 NS	0.34 *	0.49 **
<b>GFD</b>	-0.62 **	-0.70 **	0.47 **	1 **	-0.23 NS	0.32 NS	0.01 NS	0.53 **	0.43 **	-0.42 **	-0.25 NS	0.61 **	0.39 *	0.10 NS	0.34 *
<b>ATN</b>	-0.03 NS	-0.03 NS	0.13 NS	0.12 NS	1 **	-0.13 NS	-0.47 **	0.50 **	0.54 **	-0.42 **	-0.38 *	-0.08 NS	-0.62 **	0.09 NS	0.38 *
<b>AFL</b>	-0.30 **	-0.24 **	0.04 NS	0.25 **	0.20 *	1 **	0.79 **	-0.44 **	0.14 NS	1.06 **	0.33 *	0.30 NS	-0.02 NS	0.32 NS	0.53 **
<b>AFW</b>	-0.07 NS	0.01NS	0.04 NS	0.01 NS	-0.04 NS	0.50 **	1 **	-0.23 NS	0.27 NS	-0.01NS	0.47 **	0.43 **	0.13 NS	0.11 NS	0.32 NS
<b>APH</b>	0.10 NS	0.03 NS	0.26 **	0.16 NS	0.13 NS	0.04 NS	0.08 NS	1 **	0.80 **	-0.64 **	0.47 **	0.73 **	-0.05 NS	-0.37 *	0.65 **
<b>BYL</b>	-0.24 *	-0.16 NS	0.26 **	0.34 **	0.30 **	0.21 *	0.17 NS	0.44 **	1 **	-0.26 NS	0.18 NS	0.85 **	0.36 *	-0.21 NS	0.95 **
<b>HI</b>	0.06 NS	0.06 NS	-0.0 NS	-0.11 NS	-0.05 NS	9e-04 NS	-0.06 NS	-0.17 NS	-0.47 **	1 **	0.46 **	-0.63 **	-0.06 NS	0.53 **	0.03 NS
<b>ASL</b>	0.20 *	0.23 *	0.21 *	-0.05 NS	0.05 NS	0.07 NS	0.29 **	0.38 **	0.12 NS	0.02NS	1 **	0.04 NS	-0.31 NS	-0.06 NS	0.31 NS
<b>SPS</b>	-0.11 NS	-0.10 NS	0.25 **	0.28 **	0.02 NS	0.08 NS	0.25 **	0.32 **	0.31 **	-0.15 NS	0.13 NS	1 **	-0.08 NS	-0.43 **	0.72 **
<b>CC</b>	0.01 NS	9e-04 NS	0.15 NS	0.11 NS	-0.09 NS	0.08 NS	0.08 NS	-0.10 NS	-0.04 NS	0.03 NS	-0.08 NS	0.05 NS	1**	0.01 NS	0.35 *
<b>TW</b>	0.08 NS	-0.08 NS	0.18 *	0.05 NS	0.08 NS	0.03 NS	0.04 NS	-0.16 NS	-0.09 NS	0.18 NS	0.13 NS	-0.24 *	0.03 NS	1**	0.35 *
<b>YL</b>	-0.24 **	-0.16 NS	0.25**	0.33 **	0.32 **	0.22 *	0.16 NS	0.38 **	0.86 **	0.09 NS	0.13 NS	0.38 **	-0.03 NS	-0.03 NS	1**

Fig3: Pearsons Correlation

Fig: Pearsons Correlation



### 3.2. Path coefficient analysis

Phenotypic and Genotypic path coefficient analysis is presented in Table 4&5. The direct impact of characters on grain yield revealed that their relationships significantly contributed to the final grain yield, making them pivotal factors in enhancing grain production. Genotypic correlation coefficients were partitioned using the path analysis method to determine the direct and indirect effects of yield contributing traits towards grain yield. Path analysis (Table.4 & 5) revealed that the highest positive direct effect and genotypic correlation with grain yield were obtained by biological yield, harvest index, days of anthesis, date of maturity, grain filling period, seeds per spike, days of heading, flag leaf length, similar results reported by (Jouyban *et al.*, 2015 & Solomon, 2013)<sup>[9,16]</sup>. Plant height, thousands of seeds weight, spike length, flag leaf width, adequate tiller number and chlorophyll content had a negative direct effect with significant genetic correlation with grain yield.

On the other hand, days to heading had a positive indirect effect with anthesis, maturity, flag leaf width, harvest index, chlorophyll content and adverse indirect effects with grain filling period, tiller number, flag leaf length, plant height, biological yield, spike length, thousands seeds weight. The residual effect of the present study was -0.0055, indicating that about 98 per

cent of the variability in grain yield might be contributed by these 16 yield-contributing traits studied in the path analysis. This gives the impression that some other minor characters than those involved in the present study also contributed to the variability of grain yield.

UNDER PEER REVIEW

**Table 4: Genotypic Path analysis**

Traits	DTH	DTA	DTM	GFD	ATN	AFL	AFW	APH	BYL	HI	ASL	SPS	CC	TW
<b>DTH</b>	<b>0.074</b>	0.258	0.018	-0.111	-0.028	-0.088	0.029	-0.079	-0.034	0.003	-0.100	-0.042	0.069	-0.038
<b>DTA</b>	0.071	<b>0.266</b>	0.030	-0.106	-0.036	-0.034	-0.004	-0.040	0.058	0.049	-0.141	-0.065	0.052	-0.057
<b>DTM</b>	0.007	0.046	<b>0.173</b>	0.104	0.011	0.019	-0.007	-0.272	0.705	-0.133	-0.072	0.025	-0.018	-0.097
<b>GFD</b>	-0.050	-0.172	0.111	<b>0.163</b>	0.037	0.041	-0.003	-0.179	0.501	-0.141	0.054	0.070	-0.055	-0.030
<b>ATN</b>	0.013	0.062	-0.012	-0.039	<b>-0.155</b>	-0.017	0.069	-0.169	0.638	-0.143	0.083	-0.010	0.087	-0.027
<b>AFL</b>	-0.051	-0.071	0.026	0.053	0.021	<b>0.126</b>	0.115	0.148	0.172	0.354	-0.073	0.034	0.003	-0.091
<b>AFW</b>	-0.015	0.007	0.008	0.003	0.074	0.100	<b>-0.145</b>	0.080	0.318	-0.006	-0.101	0.049	-0.019	-0.031
<b>APH</b>	0.017	0.032	0.140	0.087	-0.078	-0.055	0.034	<b>-0.337</b>	0.938	-0.215	-0.103	0.084	0.007	0.104
<b>BYL</b>	-0.002	0.013	0.105	0.070	-0.085	0.019	-0.040	-0.272	<b>1.163</b>	-0.087	-0.039	0.098	-0.050	0.060
<b>HI</b>	0.001	0.039	-0.069	-0.069	0.067	0.134	0.002	0.218	-0.307	<b>0.332</b>	-0.100	-0.073	0.010	-0.150
<b>ASL</b>	0.034	0.174	0.058	-0.041	0.060	0.043	-0.068	-0.161	0.213	0.154	<b>-0.215</b>	0.005	0.044	0.018
<b>SPS</b>	-0.027	-0.152	0.038	0.100	0.014	0.038	-0.063	-0.249	1.000	-0.211	-0.009	<b>0.114</b>	0.012	0.122
<b>CC</b>	-0.036	-0.100	0.022	0.064	0.097	-0.003	-0.020	0.017	0.421	-0.023	0.068	-0.010	<b>-0.139</b>	-0.004
<b>TW</b>	0.010	0.054	0.060	0.018	-0.015	0.041	-0.016	0.125	-0.247	0.177	0.014	-0.050	-0.002	<b>-0.281</b>

**Table 5: Phenotypic path analysis**

Traits	DTH	DTA	DTM	GFD	ATN	AFL	AFW	APH	BYL	HI	ASL	SPS	CC	TW
<b>DTH</b>	<b>-0.012</b>	-0.167	0.043	0.123	0.000	0.010	-0.002	-0.002	-0.267	0.035	-0.003	-0.003	0.000	-0.003
<b>DTA</b>	-0.010	<b>-0.186</b>	0.046	0.139	0.000	0.008	0.001	-0.001	-0.185	0.033	-0.003	-0.003	0.000	-0.003
<b>DTM</b>	-0.003	-0.055	<b>0.155</b>	-0.093	0.001	-0.001	0.001	-0.006	0.294	-0.043	-0.003	0.006	0.002	-0.006
<b>GFD</b>	0.007	0.131	0.073	<b>-0.197</b>	0.001	-0.009	0.000	-0.004	0.388	-0.062	0.001	0.007	0.001	-0.002
<b>ATN</b>	0.000	0.006	0.021	-0.025	<b>0.005</b>	-0.007	-0.001	-0.003	0.334	-0.003	-0.001	0.001	-0.001	-0.003
<b>AFL</b>	0.004	0.046	0.006	-0.050	0.001	<b>-0.034</b>	0.014	-0.001	0.234	0.000	-0.001	0.002	0.001	-0.001
<b>AFW</b>	0.001	-0.004	0.007	-0.003	0.000	-0.017	<b>0.027</b>	-0.002	0.196	-0.036	-0.004	0.006	0.000	-0.002
<b>APH</b>	-0.001	-0.006	0.042	-0.033	0.001	-0.001	0.002	<b>-0.021</b>	0.497	-0.097	-0.005	0.008	-0.001	0.005
<b>BYL</b>	0.003	0.031	0.041	-0.069	0.002	-0.007	0.005	-0.009	<b>1.111</b>	-0.255	-0.002	0.010	-0.001	0.003
<b>HI</b>	-0.001	-0.011	-0.012	0.023	0.000	0.000	-0.002	0.004	-0.522	<b>0.541</b>	0.000	-0.004	0.000	-0.006
<b>ASL</b>	-0.002	-0.044	0.034	0.011	0.000	-0.002	0.008	-0.008	0.136	0.013	<b>-0.013</b>	0.003	0.000	0.000
<b>SPS</b>	0.001	0.020	0.040	-0.057	0.000	-0.003	0.007	-0.007	0.437	-0.083	-0.002	<b>0.025</b>	0.001	0.008
<b>CC</b>	0.000	0.000	0.024	-0.023	0.000	-0.003	0.000	0.002	-0.046	0.002	0.000	0.001	<b>0.012</b>	-0.001
<b>TW</b>	-0.001	-0.016	0.029	-0.012	0.000	-0.001	0.001	0.003	-0.101	0.099	0.000	-0.006	0.000	<b>-0.032</b>

## Conclusion

The current investigation, titled "Genetic Variability, Heritability, Correlation, and Path Coefficient Studies for Yield and Yield-Related Components of Selected Barley Cultivars and Crosses (*Hordeum vulgare* L.)," aimed to conduct phenotypic characterization and assess divergence among various barley genotypes. In each variety, five plants were selected for various morphological observations. Phenotypic data for various morphological traits were analyzed with the help of some statistical tools like Mean, Range (Minimum and maximum), Standard Error of mean, Critical variance (5%), Critical difference (1%), Environmental variance, Genotypic variance, Phenotypic variance, Environmental coefficient of variance, Genotypic coefficient of variance, Phenotypic coefficient of variance, Heritability (Broad sense), Genetic advance. A Higher genotypic coefficient of variation was found in no. Biological yield, grains yield, followed by seed per spike, average flag leaf width, thousand grains yield and higher phenotypic coefficient variation were found in biological yield and followed by grains yield, average flag leaf width, leaf area, and germination percentage, respectively. Genetic advance is highest in biological yield followed by seeds per spike, thousand-grain yield and grains yield (Table.2). Therefore, a correlation study revealed that days to anthesis, average spike length showed a strong positive association with biological yield, grain yield per genotype and grain filling period, average flag leaf length, seeds per spike and chlorophyll content showed a negative association with many traits. Therefore, the obtained results indicate the presence of sufficient genetic variability for the studied traits, showing that genotypes are suitable for breeding purposes.

## Acknowledgements

I would like to express my sincere gratitude to *CCS University, Meerut* for providing me with the resources and support necessary to complete this research. Special thanks to *Prof. Rahul Kumar, Head of the department of genetics and plant breeding* for his invaluable guidance, encouragement, and insightful feedback throughout the course of this project. His expertise and mentorship have been instrumental in shaping this work. I am deeply grateful for the opportunity to learn under his supervision.

## Abbreviations

**DTH:** Days to heading, **DTA:** Date of 50% anthesis, **DTM:** Date of maturity, **GFD:** Grain filling period, **APH:** Average plant height, **FL:** Flag leaf length, **FW:** Flag leaf width,

**SL:**spike length, **ATN:** Average tiller number, **NG:** No. of grains per spike, **BY:** biological yield, **GER:** Germination percentage, **CC:** Chlorophyll content, **GY:** grain yield, **TGW:** thousand-grain yield, **HI:** Harvest index, **&:** and, **%:** percent, **SEm:** Standard error of mean, **CV:** Critical variance (5%), **CD:** Critical difference (1%), **GCV:** Genotypic coefficient of variance, **PCV:** Phenotypic coefficient of variance, **Heri.(BS):** Heritability (Broad sense), **GA:** Genetic advance.

## References

- [1] Addisu, F., & Shumet, T. (2015). Variability, heritability and genetic advance for some yield and yield related traits in barley (*Hordeum vulgare* L.) landraces in Ethiopia. *International Journal of Plant Breeding and Genetics*, 9(2), 68-76.
- [2] Allard, R. W., Jain, S. K., & Workman, P. L. (1968). The genetics of inbreeding populations. *Advances in genetics*, 14, 55-131.
- [3] Bothmer, R. von, and N Jacobson. 1985. Origin, taxonomy, and related species. In *Barley*, ed. D. C. Rasmusson, 19-56. Madison, Wis.
- [4] Bourne, T. F., & Poehlman, J. M. (1987). Evaluation of barley yellow dwarf tolerance from plant traits in two winter× spring barley crosses. *Euphytica*, 36(2), 585-589.
- [5] Breeding, P., & Singh, B. D. (1985). Kalyani publishers. New delhi.
- [6] Falconer, D. S. (1996). *Introduction to quantitative genetics*. Pearson Education India.
- [7] Hill, W. G., & Mackay, T. F. (2004). DS Falconer and *Introduction to quantitative genetics*. *Genetics*, 167(4), 1529-1536.
- [8] Jain, S.K. and Allard, R.W., 1960. Population studies in predominantly self-pollinated species, I. Evidence for heterozygote advantage in a closed population of barley. *Proceedings of the National Academy of Sciences of the United States of America*, 46(10), p.1371.
- [9] Jouyban, A., Give, H. S., & Noryan, M. (2015). Relationship between agronomic and morphological traits in barley varieties under drought stress condition. *Intl. Res. J. Appl. Basic. Sci*, 9(9), 1507-1511.
- [10] Karsai, I., Hayes, P. M., Kling, J., Matus, I. A., Mészáros, K., Láng, L., & Sato, K. (2004). Genetic variation in component traits of heading date in *Hordeum vulgare* subsp. *spontaneum* accessions characterized in controlled environments. *Crop science*, 44(5), 1622-1632.

- [11] Karthikeyan, P., Anbuselvam, Y., Elangaimannan, R., & Venkatesan, M. (2010). Variability and heritability studies in rice (*Oryza sativa* L.) under coastal salinity. *Electronic Journal of Plant Breeding*, 1(2), 196-198.
- [12] Narwal, S., Kumar, D., Sheoran, S., Verma, R. P. S., & Gupta, R. K. (2017). Hulless barley as a promising source to improve the nutritional quality of wheat products. *Journal of food science and technology*, 54(9), 2638-2644.
- [13] Poehlman, J. M., & Poehlman, J. M. (1987). Breeding barley and oats. *Breeding Field Crops*, 378-420.
- [14] Riaz, R., & Chowdhry, M. A. (2003). Genetic analysis of some economic traits of wheat under drought condition. *Asian Journal of Plant Sciences*.
- [15] Singh, V. P., Man, S., & Lal, J. P. (2000). Gamma ray and EMS induced genetic variability for quantitative traits in urdbean (*Vigna mungo* L. Hepper). *Indian Journal of Genetics & Plant Breeding*, 60(1), 89-96.
- [16] Sivasubramaniam, S., & Madhava Menon, P. (1973). Genotypic and phenotypic variability in rice.
- [17] Solomon, G. (2013). Correlation and path analysis in yield and yield components in spring bread wheat (*Triticum aestivum* L.) genotypes under irrigated condition in Southern India. *African Journal of Agricultural Research*, 8(24), 3186-3192.
- [18] Verma, R. P. S., Kharub, A. S., Sarkar, B., & Kumar, D. (2011). Barley: A crop for changing climate in India. *Progressive Agriculture*, 11(conf), 63-73.
- [19] Von Bothmer, R., & Seberg, O. (1995). Strategies for the collecting of wild species. *Collecting plant genetic diversity, technical guidelines*. CAB International, Wallingford, United Kingdom, 93-111.
- [20] Von Bothmer, R., Giles, B. E., & Jacobsen, N. (1986). Crosses and genome relationship in the *Hordeum patagonicum* group. *Genetica*, 71(1), 75-80.
- [21] Yadav, S. K., Singh, A. K., Pandey, P., & Singh, S. (2015). Genetic variability and direct selection criterion for seed yield in segregating generations of barley (*Hordeum vulgare* L.). *American Journal of Plant Sciences*, 6(09), 1543.

UNDER PEER REVIEW