

## Original Research Article

### **Chemical and physical characterization of vermicompost produced from organic substrates.**

#### **ABSTRACT**

The present research work was based on the evaluation of some physical and chemical properties of humus using *Eisenia foetida* for the transformation of substrates (bovine and ovine manure and cocoa shells). It was established under a completely randomized experimental design, three substrates were studied together with their combinations without repeating them, giving a total of six substrates with ~~4~~ four repetitions each, obtaining 24 experimental units. The variables evaluated were major elements (N, P, K, S, Ca and Mg), as well as the carbon-nitrogen ratio, pH, organic matter, granulometry and electrical conductivity. The results found in this research demonstrate that the quality and quantity of the fertilizers are affected by the different substrates evaluated, with differences being observed between treatments due to the materials that make up each treatment, where the type of manure used has a direct influence. The choice of materials and their pre-composting processes, depending on the material to be used, must be adequate to feed the earthworms, since an unanalyzed decision can increase the mortality of the earthworms by 100%. It is important to point out that the origin, age, and storage of the materials used influence the mineral composition. From the results obtained in this study, it is understood that the mineral composition and some physical parameters of the worm humus will be generally similar to each other.

**Keywords:** agroecological alternatives, organic fertilizers, humus.

#### **INTRODUCTION**

Organic agriculture uses a great variety of technological options with the aim of reducing and retrieving production costs, protecting health, improving the quality of life and the quality of the environment, ~~while and~~ intensifying biological interactions and beneficial natural **processes** [1].

Residual organic materials, such as those coming from harvest and livestock activities, usually end up in the environment, saturating food chains, however, they may have a better destiny if they are transformed into **composts, which** are potentially reliable for use as ~~a~~ mean for ~~the~~ plant growth [2].

The harnessing of ~~the~~ agro-industrial waste in the production of agricultural substrates **is** an important alternative for the recycling of these materials, in addition to ~~the~~ **their** agronomic, social, economic, and ecological importance, contributing in a positive way to the increase ~~of the~~ **in** production and the improvement of crop **quality** [3].

These wastes, after going through the vermicomposting process, present ideal characteristics to be used as **substrates** [4]. Vermicomposting is defined as the degradation and biological stabilization of organic matter, after the digestion of organic waste by earthworms, ~~being~~ **with** the species *Eisenia foetida* ~~being~~ the most **used** [5]. The product of vermicomposting ~~gives is~~ an organic fertilizer obtained from plant waste or animal manure, previously stabilized and neutral [6].

In the state of Chiapas, exist a large amount of organic matter left over from agricultural activities, such as cocoa husks and livestock activities related to cattle, horse, and sheep production, which are sometimes not used properly and can affect the environment because they are carried directly to drainage systems and bodies of water.

In this context, the present study proposes the use of agro-industrial residues from cocoa husks mixed with different types of manure to obtain organic substrates through vermicomposting, using the Californian red earthworm (*Eisenia foetida*).

## MATERIALS AND METHODS

The research was carried out in the municipality of Huehuetán, in the state of Chiapas. The experimental site is geographically located at  $15^{\circ}22'40,96''$  LN and  $-92^{\circ}22'40,96''$  LO with an altitude of 65 masl, as shown in Figure 1.



Figure 1. Geographic location of the study area.

The type of climate that predominates in Huehuetán, Chiapas, according to García [7], has a warm humid climate with summer rains that corresponds to Aw (w) ig. The National Meteorological Service (NMS) with its climatological normals, mentions that the annual rainfall in Huehuetán is 2420.5 mm and has an average maximum temperature of  $34.6^{\circ}\text{C}$ , average of  $28.2^{\circ}\text{C}$  and a minimum of  $21.8^{\circ}\text{C}$ .

### Experimental Design

A completely randomized experimental design was employed, using three substrates and their combinations at 50%, resulting in six treatments. Considering four replications per treatment, a total of 24 experimental units were obtained. To differentiate one treatment from the others, a letter with its respective number was assigned for the total number of replications for each treatment, presented as follows:

T1- Bovine manure (100%)

T2- Sheep manure (100%)

T3- Cocoa husk (100%)

T4- Bovine manure (50%) / Sheep manure (50%)

T5- Bovine manure (50%) / Cocoa husk (50%)

T6- Sheep manure (50%) / Cocoa husk (50%)

### **Experimental Materials**

The vermicomposting stage was carried out in commercial plastic grids (37.79 cm wide x 54.48 cm long x 32.5 cm high), with a total volume of 0.066 m<sup>3</sup>, of which 0.03 m<sup>3</sup> were used with fresh material from each of the treatments. The grids were covered with small-pore wire mesh to prevent the entry of animals and insects and to favor aeration.

The materials used (bovine manure, sheep manure and cocoa husks) were pre composted before inoculation with the worm, to provide adequate conditions to the substrate for the adaptation of the worm. Once the grids were inoculated, they were watered daily to maintain the substrate humidity at 80%.

The earthworm *Eisenia foetida* also called "red Californian" was used, which was placed at a density of 100 adults per 0.03 m<sup>3</sup> of fresh material. The worms used were mature and developed individuals, which presented the clitellar structure formed. After three months, the humus was recovered for its evaluation and characterization through chemical and physical properties.

### **Chemical Analysis**

Chemical analyses were performed according to the Mexican standard for vermicompost [8], where the percentage of organic matter (OM), nitrogen (N), phosphorus (P), potassium (K), magnesium (Mg), sulfur (S), calcium (Ca), C/N ratio, pH and electrical conductivity (EC) were determined.

The pH was determined with a potentiometer on approximately 50 g of saturated paste of the material and after 1 h of rest. The saturated paste was then filtered, the liquid was collected, and the electrical conductivity (EC) was measured, the units used were  $\text{dSm}^{-1}$ . The calculation of organic matter (OM) and the carbon-nitrogen ratio (R:C/N) were carried out with the determination of organic carbon using the Walkey-Black method. The major elements N, P, K, S, Ca and Mg were determined after total digestion of the compost, a process by which all the organic components are mineralized. These elements were then determined by atomic absorption photometric spectrometry and colorimetry, the latter for P. The N was determined by the Kjeldahl method. All these analyses were carried out in the soil laboratory of the Faculty of Agricultural Sciences (FCA), Campus IV, of the Autonomous University of Chiapas (UNACH).

### **Physical analysis**

Physical analyses were carried out according to the Mexican standard [8]. The EC was measured at a fertilizer water ratio (1:5 w/v) with a potentiometer (Orion Star model A211, Iowa, USA). The mean particle diameter (Mpd) was determined by granulometric analysis, using the sieving method for the ASTM  $\sqrt{2}$  sieve series.

### **Statistical Analysis**

The data obtained was processed using SAS software version 9.0. To establish the statistical difference between the variables, an analysis of variance (ANOVA) was performed and in cases statistical differences were found, they were compared with Tukey's test ( $p \leq 0.05$ ).

## RESULTS AND DISCUSSION

### Characteristics of the obtained vermicompost.

#### Phosphorus (P), potassium (K), calcium (Ca), sulfur (S) and magnesium (Mg).

Table 1 shows, apart from N, the total element contents for the different vermicomposts evaluated. The results show that the quality and quantity of the composts are affected by the different materials used.

**Table 1.** Mean values of phosphorus (P), potassium (K), calcium (Ca), sulfur (S) and magnesium (Mg) in vermicompost produced with *E. foetida*.

Treatment	P (%)	K (%)	Ca (%)	S (%)	Mg (%)
1	0.323 <sup>b</sup>	0.400 <sup>d</sup>	0.413 <sup>c</sup>	1.460 <sup>ba</sup>	0.120 <sup>ba</sup>
2	0.486 <sup>a</sup>	1.29 <sup>cb</sup>	0.696 <sup>a</sup>	1.776 <sup>a</sup>	0.166 <sup>a</sup>
3	0.043 <sup>d</sup>	2.723 <sup>a</sup>	0.593 <sup>ba</sup>	0.853 <sup>d</sup>	0.096 <sup>b</sup>
4	0.373 <sup>b</sup>	0.623 <sup>d</sup>	0.480 <sup>bc</sup>	1.273 <sup>bc</sup>	0.146 <sup>ba</sup>
5	0.150 <sup>c</sup>	0.650 <sup>cd</sup>	0.420 <sup>c</sup>	0.973 <sup>dc</sup>	0.126 <sup>ba</sup>
6	0.366 <sup>b</sup>	1.816 <sup>b</sup>	0.660 <sup>a</sup>	1.360 <sup>b</sup>	0.130 <sup>ba</sup>
CV	12.46%	18.71%	11.1%	10.29%	15.35%
ESx	0.001	0.054	0.003	0.017	0.000

\* Values with equal letters per column do not differ statistically ( $P \leq 0.05$ ). Treatment: 1: bovine manure, 2: sheep manure, 3: cocoa husk, 4: bovine manure (50%)/sheep manure (50%), 5: bovine manure (50%)/cocoa husk (50%), 6: sheep manure (50%)/cocoa husk (50%).

In the case of P, treatment T2 (100% sheep manure) stands out with a value of 0.486 %, differing statistically from the rest of the treatments. This result corresponds to that cited by Rincones [9], who propose adequate values of 0.4 % or higher. Similar results

were obtained by Srivastava [10], who obtained the highest P value in vermicompost from manure.

Treatment T3, with the highest percentage of cocoa husk, has the lowest value among treatments (0.043%), because of its vegetable origin and has no nutritional value. This behavior could have influenced the values observed in treatment T5 (the second lowest value) with in the presence of 50% cocoa husk. Similar results were reported by David-Santoya [11], who obtained the lowest phosphorus values (0.2%) with the highest combination of cocoa husk in their treatments.

In relation to K (Table 1), the value was particularly high in the cocoa husk material (T3), due to the high amounts that are naturally absorbed and translocated by this crop from the soil and that remain in these wastes. Studies obtained by Rojas-Molina [12] and related to the rate of decomposition and release of nutrients in cocoa residues, show differences in the results depending on the part of the plant being evaluated as a source; in this case, the fruit is the major sink for K.

In the case of Ca, there were significant differences between the materials, being higher in sheep manure (0.696 %) and its combination with cocoa husk (0.660 %). However, these values are below those obtained by Sharma and Garg [13], who report concentrations of 2.3 %. To increase the total calcium contents in the treatments, it is advisable to place crushed eggshells, since Antoni-Huanca [14] mentions that the external layers of eggshells can contain up to 95 % calcium carbonate by weight and when incorporated in vermicompost, these will increase their Ca concentrations.

In the Mg, the differences between the materials were minor, with the highest value in sheep manure (T2), slightly below the results found by Hernández-López [15] with a result of 0.28 % magnesium in horse manure. With respect to sulfur, sheep manure (T2) was the one that presented the highest percentage (1.776%), being below the results

obtained by Capulin-Grande [16] who reported a concentration of 4.8 % of S in liquid bovine manure.

### ***Hydrogen potential (pH) and electrical conductivity (EC)***

As shown in Table 2, in five of the six treatments, the pH values were greater than 7. The highest pH value was for the combination of sheep manure and cocoa husk (9.37), while the lowest was for the bovine manure (6.69). The alkaline pH of vermicompost can be attributed to the initial decomposition of organic matter, the formation of ammonium ions, and the generation of organic compounds with an alkaline reaction [17]. Huaccha [17] reports that, during vermicomposting, alkalization of the medium takes place due to the generation of ammonium and degradation of organic acids by the activity of aerobic organisms.

According to Ramírez-Gerardo [1], the pH values recorded (6.69 to 9.37) in the vermicompost indicate that they went through a probably insufficient aerobic degradation process, since ideally, the expected pH after a composting process should be neutral to slightly basic, so it is possible that the processes for composting for this study are relatively incomplete, probably because the final cooling stage, where the mesophilic microbial activity restarts [18], does not occur or is very short.

However, it should be remembered that the function of Morren's glands within the worm morphology is to secrete calcium carbonate and produce alkaline digestion, so slightly alkaline pH values are to be expected in different worm humus [13].

**Table 2.** Mean values of pH and electrical conductivity (EC) in vermicompost produced with *E. foetida*.

Treatments	pH	EC (dSm <sup>-1</sup> )
------------	----	-------------------------

1	6.69 <sup>d</sup>	4.296 <sup>b</sup>
2	7.90 <sup>b</sup>	6.920 <sup>a*</sup>
3	7.42 <sup>c</sup>	2.656 <sup>c</sup>
4	7.27 <sup>c</sup>	6.096 <sup>a</sup>
5	8.89 <sup>b</sup>	3.456 <sup>cb</sup>
6	9.37 <sup>a*</sup>	9.370 <sup>a</sup>
<b>CV</b>	<b>1.485</b>	<b>10.779</b>
<b>ESx</b>	<b>0.012</b>	<b>0.249</b>

\* Values with equal letters per column do not differ statistically ( $P \leq 0.05$ ). Treatment: 1: bovine manure, 2: sheep manure, 3: cocoa husk, 4: bovine manure (50%)/sheep manure (50%), 5: bovine manure (50%)/cocoa husk (50%), 6: sheep manure (50%)/cocoa husk (50%).

The means of the electrical conductivity values between treatments showed significant differences. The ranges varied from 2.65 to 9.37  $\text{dSm}^{-1}$ , i.e., they showed a very high variability, which explains the high coefficient of variation (Table 2). Similar results were obtained by López-Méndez [19], who observed that when manure and vegetable residue are added to the mixture, the EC of the vermicompost reaches 5 and 8  $\text{dSm}^{-1}$ , which if compared with the Mexican standard [8], it is observed indicates that most of the treatments exceed the established range by the standard, except for treatments T3 and T5, which were in the established range ( $\leq 4 \text{ dSm}^{-1}$ ). In this regard, Ramírez-Gerardo [1], mentions that mixing manure with vegetable residue causes an increase in EC attributable to the source material and the process of obtaining the organic fertilizer.

*Nitrogen (N), organic matter (OM) and carbon/nitrogen ratio (C/N).*

Regarding nitrogen (Table 3), the cocoa husk (T3) and sheep manure (T2) treatments showed the highest values, while the combination of bovine manure and sheep manure (T4) was the treatment with the lowest value. However, the Mexican standard [8], establishes that the optimum range should be 1% to 4% N, so the six treatments are within the established range.

**Table 3.** Mean values of nitrogen (N), organic matter (OM) and carbon/nitrogen ratio (C/N) in vermicompost produced with *E. foetida*.

Treatments	N (%)	OM (%)	C/N
1	1.576 <sup>bc</sup>	30.358 <sup>bc</sup>	11.17 <sup>a*</sup>
2	2.060 <sup>ba</sup>	25.570 <sup>dc</sup>	7.208 <sup>b</sup>
3	2.116 <sup>a*</sup>	37.089 <sup>a*</sup>	10.163 <sup>a</sup>
4	1.436 <sup>c</sup>	25.946 <sup>dc</sup>	10.771 <sup>a</sup>
5	1.94 <sup>bac</sup>	33.284 <sup>ba</sup>	10.005 <sup>a</sup>
6	1.99 <sup>ba</sup>	21.550 <sup>d</sup>	6.28 <sup>b</sup>
<b>CV</b>	<b>10.4%</b>	<b>7.766</b>	<b>10.406%</b>
<b>ESx</b>	<b>0.037</b>	<b>5.06</b>	<b>0.929</b>

\* Values with equal letters per column do not differ statistically ( $P \leq 0.05$ ). Treatment: 1: bovine manure, 2: sheep manure, 3: cocoa husk, 4: bovine manure (50%)/sheep manure (50%), 5: bovine manure (50%)/cocoa husk (50%), 6: sheep manure (50%)/cocoa husk (50%).

There was a significant difference between T6 (sheep manure and cocoa husk) and the others, since it had a lower organic matter (OM) content. The range of the means was 21.55 to 37.09% (Table 3), which, compared to the Mexican standard [8], indicates that vermicompost should contain 20% to 50% OM. This shows that the OM contents of the six treatments are within the optimum range.


The OM, besides having a direct influence on soil fertility, is an important factor in the presence and availability of micronutrients [20]. Therefore, in treatment T3, which presented a higher organic matter content, micronutrients are probably more readily available. This translates into a richer compost in terms of humified organic matter content and with more available nutrients for the plant to take advantage of [2].

Regarding the C/N ratio (Table 3), treatment T1 (bovine manure) was the one that showed the highest value, possibly due to the high amounts of remaining fiber that comes come from the animals' diet and that, in a certain way, remains remain in good quantities in the raw material used. Treatments T3, T4 and T5 did not differ statistically from treatment T1. Based on the Mexican Standard [8], all treatments have an adequate C/N value ( $< 20$ ).

Authors such as Gamarra-Lezcano [21] and Raza [22] mention that C/N ratios between 10 and 25, are indicative of adequate mineralization by the microorganisms present in the materials, which implies that with low C/N values, the microorganisms will be more efficient in the decomposition of organic matter.

### **Granulometry**

The substrates analyzed showed statistically significant differences in relation to particle size (Figure 2). Treatments (T1) and (T2) presented the highest percentage of particles smaller than 0.25 mm. All substrates presented the highest percentage of particles with sizes between 0.25 and 2 mm, which is considered to have a fine particle size that facilitates the supply of nutrients to the soil and to the plants. Treatment T6 was the substrate with the highest percentage of particles larger than 2 mm, in addition to the lowest proportion of particles smaller than 0.25 mm. According to González-Cueto [23], in coarser materials, the particle size can be adjusted by grinding processes.



**Figure 2.** Granulometry of vermicompost produced with *E. foetida*. Treatments with equal letters per column do not differ statistically ( $P \leq 0.05$ ). Treatment: 1: bovine manure, 2: sheep manure, 3: cocoa husk, 4: bovine manure (50%)/sheep manure (50%), 5: bovine manure (50%)/cocoa husk (50%), 6: sheep manure (50%)/cocoa husk (50%).

According to Abad [24], substrates must have a granulometry of 0.25 to 2.5 mm to be used in the field. The evaluation of eleven types of materials with potential use as substrates of plant and mineral origin carried out by Camacho [25], reported that more than 65% of the particles are within the range of 0.25 to 2.5 mm, coinciding with the results obtained in this work, which allows establishing that the substrates obtained can be used for the desired purposes.

## CONCLUSIONS

Vermicomposting is a sustainable option to use organic wastes generated in the region for the generation of vermicompost, which can be used as organic fertilizer to increase soil fertility.

The concentration of nutrients in vermicompost varies according to the components of the substrate; however, the values obtained in the study are similar to those reported in the specialized literature.

The presence of *Eisenia fetida* in the vermicomposting process ensures the production of a safe and sustainable product.

## RECOMMENDATIONS

It is recommended that the humus obtained in this study be applied in a crop to analyze its performance and check its effectiveness.

Experiment with other types of substrates to compare their nutritional value in humus.

## References

1. Ramirez-Gerardo, M. G., Vázquez-Villegas, S., Méndez-Gómez, G. I., & Mejía-Carranza, J. (2021). Characterization of organic fertilizers applied to flower crops in the south of the State of Mexico. UAT Science, 16(1), 150-161. <https://doi.org/10.29059/cienciauat.v16i1.1518>
2. Xavier, A., Delgado, G., Carlos, J., Delgado, G., Alberto, C., & Navarrete, D. (2022). Analysis of the impact on these waste sludge treatment community by means of wetlands and vermicomposting and the production of fertilizer: systematic review. Interdisciplinary Scientific Journal of Research and Knowledge, 12(1), 1-22.
3. Correa, B. A., Parreira, M. C., Martins, J. S., Ribeiro, R. C., & Silva, E. M. (2019). Reuse of regional agro-industrial organic waste from the Amazon Tocantina as

- alternative substrates in the production of lettuce seedlings. *Brazilian Journal of Sustainable Agriculture*, 9(1), 97-104.
4. Blanco-Villacorta, M. W. (2023). Vermicomposting: an alternative to enhance urban agriculture. *Research Journal and Agricultural and Natural Resources Innovation*, 10(1), 90-103. <https://doi.org/10.53287/siha3115kw72x>
5. Lima, M. V. G., dos Santos Filho, C. A., Ferreira, J. V. V., de Souza, K. G., Shockness, L. D. S. F., & Bento, G. F. (2019). Vermicomposts as substrates in the performance of lettuce and arugula seedlings. *Green Journal of Agroecology and Sustainable Development*, 14(3), 374-381. <http://doi:10.18378/rvads.v14i3.6499>
6. Alvarez, J. M., Pasian, C., Lal, R., López, R., & Fernández, M. (2019). Vermicompost and biochar substrates can reduce nutrient leachates on containerized ornamental plant production. *Brazilian Horticulture*, 37(1), 11-25. <http://dx.doi.org/10.1590/s0102-053620190107>
7. García, E. 1973. Modifications to the Köppen climate classification system. *Institute of Geography*. UNAM. Mexico, D. F. p. 91.
8. Official Mexican Standard. 2008. Worm humus (worm compost)-specifications and testing methods. NMX-FF-109-SCFI-2008. Official Gazette of the Federation. Mexico, D. F. 24 p.
9. Rincones, P., Zapata, J., Figueroa, O., & Parra, C. (2023). Evaluation of substrates on the productive parameters of the Californian red round worm (*Eisenia fetida*). *Technological information*, 34(2), 11-20. <https://dx.doi.org/10.4067/s0718-07642023000200011>
10. Srivastava, P. K., Singh, A., Kumari, S., Arora, S., Choubey, A. K., & Sinha, A. S. K. (2023). Production and characterization of sustainable vermimanure derived

- from poultry litter and rice Straw using tigerworm *Eisenia fetida*. *Bioresource Technology*, 369, 128377. <https://doi.org/10.1016/j.biortech.2022.128377>
11. David-Santoya, J. J. E., Gómez-Álvarez, R., Jarquín-Sánchez, A., & Villanueva-López, G . (2018). *Characterization of vermicomposts and their effect on the germination and growth of Capsicum chinense Jacquin. Ecosystems and agricultural resources*, 5(14), 181-190. <http://doi:10.19136/era.a5n14.1465>
12. Rojas-Molina, J., Ortiz-Cabralez, L., Escobar-Pachajoa, L. D., Rojas-Buitrago, M., & Jaimes-Suarez, Y. Y. (2021 ). *Decomposition and nutrient release in biomass by pruning of cocoa (*Theobroma cacao* L.) in Rionegro, Santander, Colombia. Mesoamerican Agronomism*, 32(3), 888-900. <http://doi:10.15517/am.v32i3.41608>
13. Sharma, K., & Garg, V. K. (2020). *Conversion of a toxic weed in to vermicompost by Eisenia fetida: Nutrient content and earthworm fecundity. Bioresource Technology Reports*, 11, 100530. <https://doi.org/10.1016/j.biteb.2020.100530>
14. Antoni Huanca, S. G. (2019). *Use of ground eggshell as a warming material in an acidic soil of Peru. Idesia (Arica)*, 37(3), 115-120. <http://dx.doi.org/10.4067/S0718-34292019000300115>
15. Hernandez-López, M., Vidaña-Martínez, S.A., & Velasquez-Chavez T.E. (2020). *Chemical and microbiological characteristics of vermicompost produced at ITSL. Science, Engineering and Development Journal Tec Lerdo*, 1(6), 35-39.
16. Capulin-Grande, J., Mohedano-Caballero, L., Sandoval-Estrada, M., & Capulin-Valencia, J. C. (2011). *Liquid bovine manure and inorganic fertilizers on tomato paste yield in a hydroponic system. Chapingo Magazine. Horticulture Series*, 17(2), 105-114. 17.

17. Huaccha, A., Fernandez, F., Quiroga, S., & Alvarez, B. (2019). Use of *Eisenia hortensis* (earthworm) in vermicomposting of organic waste. *Pakamuros Scientific Journal*, 7(2), 32-40. <https://doi.org/10.37787/aa9ncs06>
18. Montalvo, P. A. J., Dongo, L. F. O., Maraví, J. L. C., Monzón, L. A. T., Coral, M. F. C., & Figueroa, L. V. (2018). Nitrogen transformation during horse dung composting. *Clean Production Journal*, 13(22), 77-88. <https://doi.org/10.22507/pml.v13n2a9>
19. Lopez-Méndez, C., Ruelas-Ayala, R. D., Sañudo-Torres, R. R., Armenta-López, C., & Félix-Herrán, J. A. (2020). Influence of different organic substrates on the Californian red roundworm (*Eisenia foetida*). *TECHNOSCIENCE Chihuahua*, 7(2), 81-87. <https://doi.org/10.54167/tch.v7i2.662>
20. Guadarrama-Nonato, A., Mejía-Carranza, J., & Ramírez-Gerardo, M. G. (2018). Mineralization of organic matter in soils with differential management in rose cultivation. *Acta Universitaria*, 28(32), 33-41. <https://doi.org/10.15174/au.2018.1654>
21. Gamarra-Lezcano, C. C., Díaz Lezcano, M. I., Vera de Ortíz, M., Galeano, M. D. P., & Cabrera Cardús, A. J. N. (2018). Carbon- nitrogen relationship in soils of silvopastoral systems in the Paraguayan Chaco. *Mexican Journal of Forestry Sciences*, 9(46), 4-26. <https://doi.org/10.29298/rmcf.v9i46.134>
22. Raza, S. T., Wu, J., Rene, E. R., Ali, Z., & Chen, Z. (2022). Reuse of agricultural wastes, manure, and biochar as anorganic amendment: A review on its implications for vermicomposting technology. *Journal of Cleaner Production*, 360, 132200. <https://doi.org/10.1016/j.jclepro.2022.132200>
23. González-Cueto, O., Salcerio-Salaberry, R. A., Aguila-Alcantara, E., Merlán-Mesa, G., López-Bravo, E., & Machado-de Armas, J. (2023). Physical and chemical

characterization of the organomineral fertilizer Agromena - G. Journal of Agricultural Technical Sciences, 32 (3), 1-8. <https://cu-id.com/2177/v32n3e08>

24. Abad-Berjon M, Noguera-Murray P, Carrión-Benedito C. Substrates in soilless crops. In: Urrestarazu-Gavilán. Soilless cultivation. Madrid: Mundi Prensa, 2004. 113-158.

25. Camacho, O. I. M., Toro, M. C. H., & Díaz, J. S. G. (2021). Characterization of materials with potential use as substrates in soilless cultivation systems. Agricultural Science and Technology, 22 (1), e1977. [https://doi.org/10.21930/rcta.vol22\\_num1\\_art:1977](https://doi.org/10.21930/rcta.vol22_num1_art:1977)

UNDER PEER REVIEW