

Yield gap analysis and impact of Front Line Demonstrations on Productivity of Carrot in Tirap District of Arunachal Pradesh

Abstract

The carrot is an important crop in Tirap district of Arunachal Pradesh. Farmers use old varieties for cultivation so their productivity is lower than potential. So, a demonstration of high yielding variety of carrot- Pusa Rudhira was undertaken during Rabi season of 2017-18 and 2018-19 respectively in Tirap district of Arunachal Pradesh to demonstrate the scientific carrot cultivation practices. Total 20 farmers were selected from five villages during the both years of demonstration. Before the demonstration a field survey was conducted in selected villages to know in details about farmers practices of carrot. As per findings of survey, there was huge technological gap. There was potential of Pusa Rudhira variety of carrot in district 230 q/ha. Meanwhile demonstration yields were recorded at 190 and 178 q/ha while control was 128 and 116 q/ha respectively. Extension gap was 63 & 62 q/ha respectively. The benefit- Cost (B:C) ratio was 4.39 and 3:20 and 3.63 and 2.54 respectively during the both year of demonstration.

Key words: Carrot, demonstration, Pusa Rudhira, benefit cost ratio, extension gap, field survey, yield/ha

INTRODUCTION

Carrot (*Daucus carota* L.) is one of the most important cool weather root crops is grown all over the world. In India, the important carrot growing states are Uttar Pradesh, Assam, Karnataka, Andhra Pradesh, Punjab and Haryana. It is the second and third most important vegetable in England and Australia, respectively (Biratu *et al.*, 2022). It can be grown also in mild climate of the tropics. Carrot had 0.064 million ha area with the production of 0.968 million tons during 2017-18 (MoFPI and IIFPT, 2020). Being a rich source of beta carotene, a precursor of vitamin-A. Also possesses anti-oxidant properties, fix up harmful free radicals and prevent heart diseases (Dias Joao, 2014). It is generally consumed as a vegetable after cooking but also consumed fresh in salad, juice and served as an ingredient in soups and sauces. Tender roots are processed to make pickles, jam, candy and jellies. It is also used as an additive in poultry feed to intensify skin and egg yolk colour. The carrot variety Pusa Rudhira suits to this region. Front Line Demonstration was carried out at the twenty farmer's field at Tirap district of Arunachal Pradesh. Here generally in winter's period's minimum temperature goes to 8- 9°C and in summer maximum temperature reaches to 36°C. Annual rainfall is 1500-2000 mm per year.

Material and methods

In Arunachal Pradesh three types of Agro-climatic zones are prevails. The Tirap district falls under Eastern Himalayan Region (Zone II), Sub region-: Per Humid Hyper

Thermic Foothills; where hot and humidity is very common characteristics. The rains start from End of February and continue up to September. The intermediary dry spells often occurs which are very heat and humid.

Table No- 1: The weather during the research period

Month	Rainfall(mm)		Temperature °C				Relative Humidity (%)			
	2017	2018	2017		2018		2017		2018	
			Max.	Min.	Max.	Min.	M	E	M	E
April- 2017	247	186.0	34.4	12.2	35.2	13.2	82	67	81	65
May- 2017	327	117.5	35.6	14.6	36.7	15.1	86	73	89	75
June-2017	241	433.4	36.8	16.5	37.7	17.2	91	80	93	82
July-2017	347	336.6	34.2	18.4	35.4	18.9	93	83	95	81
August-2017	493	277.3	33	19.1	34.2	20.1	87	85	89	87
September-2017	371	186.2	32.3	18.8	33.6	20.1	88	84	91	86
October-2017	162	118.0	26.5	17.2	27.4	18.4	89	90	92	92
November-2017	7.6	15.4	25.1	12.3	25.9	14.1	87	82	89	88
December-2017	0	0	25.8	9.4	26.2	10.2	85	83	86	87
January-2018	12.2	12.7	25.4	8.6	26.2	9.1	85.7	88	84.9	88
February-2018	69.6	69.0	26.1	7.9	26.9	8.3	88	90	83	92
March-2018	138.2	123.0	28.7	8.8	29.1	9.2	85	81	82	83

Where Max. denotes maximum, min. denotes minimum, M denotes Morning, E denotes evening

Table 2. Package and Farmers' Practices demonstrated in Carrot Field.

Particular	Technological intervention	Existing practices	Gap
Variety	Pusa Rudhira	Local or unknown variety	Full gap
Seed rate	6 kg/ha	10 kg /ha	Full gap
Seed treatment	Seed was treated	Not treated	Full gap
Sowing method	Line sowing	Broadcasting	Full gap
Spacing	20 cm x 4 cm	Not maintained	Full gap
Application of recommended dose of manure	5 kg/ meter ²	Nil/without recommendation	Partial gap

Application of Bio fertilizer	Soil application of Azospirillum & PSB @ 2 kg/ha mix with FYM	No application	Full gap
Harvesting	Manual	Manual	No Gap

The present study was conducted in Tirap district of Arunachal Pradesh during rabi season of 2017-18 and 2018-19 respectively. The twenty farmers were selected for demonstration; from Chomoithung, Thingsa, Kheti, Lapnan and Dadam villages in Tirap district of Arunachal Pradesh. Each farmers had 0.05 ha area of plot size. The good quality seeds of carrot cv. Pusa Rudhira were procured and distributed to twenty selected farmers. All the selected farmers were trained on various aspects of carrot production technologies before conducting of demonstration through training. The field was prepared by ploughing and harrowing after kharif crops and manure at 5kg.meter² and biofertilizers were applied during last ploughing. The seeds were sown in well prepared field during second week of November. The 6 kg/ha seeds were sown at 20 cm line to line and 4 cm plant to plant distance. All the recommended practices were followed in demonstration plots while farmers practices were grown as per the farmers practices (mentioned in table 2). The data related to cost of cultivation, production, productivity, total return and net return were collected in both treatments as per schedule from all selected farmers. An average of cost of cultivation, yield and net returns of different farmers was analyzed by the given formula-

$$\text{Average} = [F_1 + F_2 + F_3 + \dots + F_n] / N \quad F = \text{Farmer}$$

$$N = \text{No. of Farmers}$$

In the present investigation, technology index was operationally defined as the technical feasibility obtained due to implementation of Frontline Demonstrations in Carrot. To find the technology gap, extension gap and technology index following formula used by [3] have been used.

$$\text{Technology Gap} = P_i (\text{Potential Yield}) - D_i (\text{Demonstration Yield})$$

$$\text{Extension Gap} = D_i (\text{Demonstration Yield}) - F_i (\text{Farmers yield})$$

$$\text{Technology index} = [(P_i (\text{Potential Yield}) - D_i (\text{Demonstration yield})) \times 100] / P_i (\text{Potential yield})$$

Results

A comparative study of productivity levels between demonstrated variety and local check is shown in Table 3. During the period of Study, it was observed that demonstration plots recorded the higher yield during the both years (190 and 178 q/ha) as compared local

check (128 and 116 q/ha). In percentage; 48 and 53 higher over local check.

As per the parameter of technology gap, there were 40 & 52 q/ha recorded; during the both years of study while the extension gap was also had huge margins e.g. 63 and 62 q/ha respectively. And the technology index was recorded 18 and 23 % respectively.

Under economic parameters all parameters had recorded good results. The cost of cultivation during the 2017 -18 was Rs.88,000 and 76,000 respectively (demonstration and control) while it was little bit increased during the next year of demonstration (2018-19) as Rs. 96,000 and 82,000 respectively (table-4). The gross return was calculated during the first year of study was Rs. 4,75,000 and 3,20,000 as compared Rs. 4,45,000 and 2,90,000, net return was Rs. 3,87,000 and 2,44,000 as compared to Rs. 3,49,000 and 2,08,000 respectively. The benefit cost ratio of demonstration plots was also recorded superior over control (4.39 and 3.20 as compared 3.63 and 2.54 respectively).

Discussion

These findings are proving that improved varieties performance reported better over the local check under similar environmental conditions. The other farmers were motivated by seeing the results in term of productivity. The yield of the frontline demonstrations and potential yield of the crop was compared to estimate the yield gaps which were further categorized into technology index (Dawuda, 2011), Adem Seid *et al.* (2019) has described that different dose of fertilizers enhanced the yield of carrot. The Biratu *et al.* (2022) and Villeneuve (2017) have proven that sowing time of carrot played crucial role for its production meanwhile Reid *et al.* (2018) and Singh *et al.* (2006) emphasized under their findings that nutrients play important role in carrot production.

The technology gap shows the difference between potential yields over demonstration yield of the technology. The potential yield of the technology (variety Pusa Rudhira) is 230 q/ha; in soil and climatic conditions of Arunachal Pradesh. The technology gap of 40 and 52 q/ha recorded during the study has raised a special concern among the development agencies like- department of Horticulture, Agriculture, KVK etc. for making a strategy for minimizing the same. Though several efforts have been done under the supervision of KVK specialist at the farmers' field; but there are existing a gap between the potential yield and demonstration yield. The better yield of demonstration plots recorded due to scientific management practices of carrot. This finding has also supported by different researchers – Chebotev, 2010, Da Silva.2008, De Carvelho. 2018, D'Hooghe. 2018, Kabir.2013 etc. The similar findings have also reported by Kassa M. *et al.*2018. Bender,I. *et al.* 2020 and Umlong R.M., 2010 have

mentioned that fertility level of soil and weather condition are important factor for carrot yield. Kharsan *et al.* (2019), Shiberu (2016) and Tagen (2021) reported that plant spacing played vital role under carrot cropping while Kovacik (2018) emphasizes that application of vermicompost during last ploughing under carrot farming was the key factor for better production. The Raginaldo (2021) demonstrated the carrot demonstration in Latin America and concluded that weed management is the pivotal factor for its good yield. The Tewari *et al.* (2019) reported that seed treatment had saved the carrot crop against different seed and soil borned diseased; which ultimately resulted better crop growth and development; by which the yield was superior as compared to without seed treatment crop of carrot.

Technology index shows the feasibility of the variety at the farmer's field. The lower value of technology index indicates the more feasibility of the particular technology. The result of study depicted in Table 3 revealed that the technology index value was 18 during first year of demonstration; which resulted into 23 after second year's demonstration at farmer's field. This proved that this type of demonstrations can play a vital role for improving the productivity of carrot in Arunachal Pradesh. Pandey *et al.* (2020) has also confirmed that under scientific package and practices farmers income can be enhanced.

During the both years of demonstration; net return was superior over control (Rs. 1,43,000 and 1,41,000 respectively) with better Benefit cost ration (4.39 and 3.63). The higher profitability and economicviability of the demonstration can enhance the farmers productivity as well as economy.

Table-3, Yield, technology gap and technology index of demonstration

Year	Fruit yield (q/ha)			Increase in productivity (%)	Technology gap (q/ha)	Extension gap (q/ha)	Technology index (%)
	Potential	Demonstration	Control				
2017- 18	230	190	128	48	40	63	18
2018-19		178	116	53	52	62	23

Table 4. Economics of Carrot

Year	Cost of Cultivation (Rs/ha)		Gross Return (Rs/ha)		Net Return (Rs/ha)		Benefit Cost ratio B:C Ratio	
	D	C	D	C	D	C	D	C
2017-18	88,000	76,000	4,75,000	3,20,000	3,87,000	2,44,000	4:39	3:20
2018-19	96,000	82,000	4,45,000	2,90,000	3,49,000	2,08,000	3.63	2.54

Where D denotes: Demonstration and C denotes: Control.

Conclusion

The results of demonstration were 48 % and 53 % higher yield over farmers practices during the both years of study respectively. This higher result came by adoption of good variety, scientific package and practices at farmer's field. Thus, the farmers of Tirap district can learn by this result for higher productivity, profitability as well as economic prosperity.

References

- Adem Seid M., Gedamu F., Mohammed D., 2019. *Effects of blended nps fertilizer rates and plant spacing on growth, yield and quality of carrot (Daucus carota L. subsp. Sativus) in Tehuledere District, Northeastern Ethiopia.* - Haramaya University, Etiopia.
- Bender I., Edesi L., Hiiesalu I., Ingver A., Kaart T., Kaldmäe H., Talve T., Tamm I., Luik A., 2020. *Organic carrot (Daucus carota L.) production has an advantage over conventional in quantity as well as in quality.* - *Agronomy*, 10(9): 1420.
- Biratu W., Molla B., Abebe H., Gebremeskel H., 2022. *Study growth, root yield and yield related character of carrot (Daucus carota) varieties under highland areas of southern tigray region, northern Ethiopia.* - *Int. J. Life Sci.*, 10(4): 317-323.
- Chebotaev, N. F., & Danko, V. N. 2010. *Carrots for fodder.* Selkhozizdat Publ., Moscow.
- Correia, N. M., & De Carvalho, A. D. (2018). Post-emergence selectivity of metribuzin to carrot. *Revista Ceres*, 65(4), 314-320.
- Cotes, B., Rämert, B., & Nilsson, U. A. (2018). A first approach to pest management strategies using trap crops in organic carrot fields. *Crop Protection*, 112, 141-148.

- Da Silva J.B.C., Vieira J.V., Lana M.M., 2008. *Processing yield of the carrot cultivar Esplanada as affected by harvest time and planting density*. - *Scientia Hort.*, 115(3): 218-222.
- Dawuda M.M., Boateng P.Y., Hemeng O.B., Nyarko G., 2011. *Growth and yield response of carrot (Daucus carota L.) to different rates of soil amendments and spacing*. - *J. Sci. Techn. (Ghana)*, 31(2).
- De Carvalho, D. F., Gomes, D. P., De Oliveira-Neto, D. H., Rouws, J. R., & De Oliveira, F. L. 2018. Carrot yield and water-use efficiency under different mulching, organic fertilization and irrigation levels. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 22(7), 445-450.
- D'Hooghe, P., Diaz, D., Brunel-Muguet, S., Dubois, J., & Kauffmann, F. 2018. Spatial variation of root yield within cultivated carrot fields is strongly impacted by plant spacing. *Scientia Horticulturae*, 241, 29-40.
- Dias, João., 2014. Nutritional and Health Benefits of Carrots and Their Seed Extracts. *Food and Nutrition Sciences*. 05. 2147-2156. 10.4236/fns.2014.522227.
- Kabir A., Ali A., Waliullah M.H., Mohay Men-Ur R.M., Rashid A., 2013. *Effect of spacing and sowing time on growth and yield of carrot (Daucus carota L.)*. - *Inter. J. Sust. Agric.*, 5(1): 29-36.
- Kassa M., Zeberhe T., Abrihaley S., Michael T.G., Hagos K., Kiros B.G., 2018. *On farm demonstration of improved carrot (Daucus carota L.) variety in EmbaAlaje district, Northern Ethiopia*. - *Inter. J. Agric. Biosci.*, 7(4): 218-221.
- Kharsan M., Nag K., Sahu D.K., Bhardwaj L.P., 2019. *To assess the effect of spacing on growth and yield of carrot (Daucus carota L.) cv. Pusa Kesar*. - *J. Pharm. Phytochem.*, 8(5S): 77-80.
- Kováčik, P., Šalamún, P., Smolen, S., Šimanský, V., & Moravčík, L. 2018. Determination of the carrot (*Daucus Carota L.*) yields parameters by vermicomposting and earthworms (*Eisenia Foetida*). *Potravinárstvo Slovak Journal of Food Sciences*, 12(1), 520-526.
- MoFPI and IIFPT, 2020. Carrot Production, Clusters, Post-Harvest Management and Value Addition in India. <http://www.iifpt.edu.in/pmfme/dpr-carrotcake.pdf>.
- Pandey, N., Rijal, S., Adhikari, H., Bhandana, B., & Adhikari, M., 2020. Production Economics and Determinants of Carrot (*Daucus carota L.*) production in Chitwan, Nepal.

International Journal of Social Sciences and Management, 7(4), 234–241.
<https://doi.org/10.3126/ijssm.v7i4.32473>.

Reginaldo L.T.R.T., Lins H.A., Sousa M.D.F., Teófilo T., Da Silva M., Mendonça V., Silva D.V., 2021. *Weed interference in carrot yield in two localized irrigation systems*. - Revista Caatinga, 34: 119-131.

Reid, J. B., Hunt, A. G., Johnstone, P. R., Searle, B. P., & Jesson, L. K., 2018. On the responses of carrots (*Daucus carota* L.) to nitrogen supply. *New Zealand Journal of Crop and Horticultural Science*, 46(4), 298-318.

Singh, S.R., 2006. Response of biofertilizers and pesticides on yield and quality of cabbage, radish and brinjal in vegetable – based rotation system. *Applied Biological Research*, 8: 33-36.

Shiberu T., Tamiru S., 2016. *Effect of intra spacing on yield and yield components of carrot (Daucus carota L. Sub Sp. Sativus)*. - Curr. Res. Agric. Sci., 3(1): 1-6.

Tegen H., Jembere M., 2021. *Influences of spacing on yield and root size of carrot (Daucus carota L.) under ridge-furrow production*. - Open Agriculture, 6(1): 826- 835.

Tewari N, Singh P, Lal C, Katiyar PK, Vaish CP ,2019. Effect of presowing seed treatment on germination, growth and yield of onion (*Allium cepa* L.). *Seed Res* ; 29:238-9.

Umlong, R.M. 2010. Growth, yield and quality of carrot (*Daucus carota* L.) as influenced by organics and lime. M.Sc. (Ag.) Thesis, Assam Agricultural University, Jorhat.

Villeneuve, F., & Latour, F., 2017. Influence of sowing time and chilling exposure on flower induction in carrot (*Daucus carota* L.). *Acta Horticulturae*, 1153, 47-54.