

## Original Research Article

### Ethnobotanical Assessment of Wild and Semi-Wild Edible Plants in Northwestern Tigray, Ethiopia

#### Abstract

*Wild and semi-wild edible plants play a vital role in food security by supplementing dietary needs. This study explored the types of wild and semi-wild edible plants consumed, their management practices, consumption modes, marketability, and major threats in the lowland and midland agroecological zones of Northwestern Tigray, Northern Ethiopia. A total of 234 informants were involved through focus group discussions, key informant interviews, household surveys, and field observations. In total, 32 wild and semi-wild edible plant species were identified: 8 in the lowland zone, 11 in the midland zone and 13 common to both. Trees were the dominant life form (65.62%), followed by tree-shrub combinations (18.75%) and shrubs (15.65%). The Malvaceae and Moraceae families were the most represented, with four species each. Fruits were the most commonly consumed part (96.6%), while leaves (3.1%) and seeds were rarely used. Consumption was primarily raw, with 74.53% of fruits in the lowland areas and 97.01% in the midland areas consumed outdoors. Overconsumption of certain fruits can lead to side effects such as abdominal bloating, nausea, heart disease, constipation, and urinary discomfort. Children were the primary collectors and sellers of these plants, using methods like plucking or long sticks for harvesting. Market analysis revealed that only 25% of these edible plants were sold locally, with *Tamarindus indica*, *Balanites aegyptiaca*, *Ziziphus spina-christi*, and *Syzygium guineense* being the most preferred. These plants provide multiple benefits, including enhancing food security, serving as energy sources, construction materials, medicines, and household utensils, as well as offering ecological services and aesthetic value. However, agricultural expansion, fuel wood collection, and overgrazing are major threats to their sustainability. Therefore, it is crucial that appropriate utilization and conservation measures are implemented by all stakeholders to ensure their long-term availability.*

**Keywords:** *Conservation, Consumption, Management, Marketability, Threat*

## 1. Introduction

Natural forests, woodlands, and wildlife have long served as primary food sources for early hunter-gatherer societies and forest-dependent communities (Seyoum et al., 2015). Among these resources, wild edible plants have provided essential sustenance since pre-agricultural times, even before the domestication of modern crops (Amente, 2017). Globally, around 3 million people rely on non-timber forest products, highlighting the significant value of wild and semi-wild edible plants (Anbessa et al., 2024). In Ethiopia alone, people consume over 300 different wild edible plant species (Asfaw, 2008). Key habitats for these wild plants include forests, wooded grasslands, scrublands, riverine environments, and farmlands across the country's diverse vegetation zones (Asfaw, 2008).

Wild and semi-wild edible plants play a critical role in food security, providing supplementary dietary needs (Teklehaymanot and Giday, 2010; Seyoum et al., 2015). They also serve as valuable sources of income for local communities (Tahir et al., 2023) and are important in traditional medicine (Guinand and Lemessa, 2001). Studies have shown that many wild edible species contain higher concentrations of vitamins and nutrients compared to domesticated agricultural plants (Asfaw, 2008). In addition to their nutritional benefits, these plants provide forage for livestock, fuelwood, and materials for construction and tools, supporting rural households (Amente, 2017; Berihun and Molla, 2017). Furthermore, they play a vital role in biodiversity conservation and environmental protection, integrating ecological systems with human livelihoods (Ruffo et al., 2002). Traditional communities worldwide have accumulated extensive indigenous knowledge regarding the use and management of wild and semi-wild edible plants, passed down through generations of interaction with their natural surroundings (John, 1993). In Ethiopia's rural areas, elders are the primary custodians of this valuable knowledge, especially in relation to the availability and management of these plants (Abebe and Ayehu, 1993).

Despite their significant roles, wild and semi-wild edible plant species in Ethiopia face numerous threats, including deforestation, forest degradation, inadequate management practices, and population pressure (Belete et al., 2021). Moreover, comprehensive studies on wild edible fruit-bearing woody species have not been adequately conducted across many regions of the country (Belete et al., 2021; Regassa et al., 2014). Therefore, identifying and documenting these species across diverse ecological settings is essential to maximize the benefits for local communities, guiding the development of management practices that ensure the conservation and sustainable use of these valuable resources (Belete et al., 2021).

In the Tigray region, similar to other areas of Ethiopia, people rely on wild and semi-wild edible trees and shrubs for income, medicinal purposes, and as a survival strategy during food shortages and throughout normal seasons. However, despite the widespread use of these plants, few studies have been conducted to document the ethnobotany of wild and semi-wild edible plants in the region (Giday and Teklehaymanot, 2023). The aim of this study was to identify and document the wild and semi-wild fruit-bearing woody plant species found in the midland and lowland areas of northwestern Tigray.

## 2. Materials and Methods

### 2.1. Study Area

The study was conducted in Northwestern Tigray, northern Ethiopia, to assess the availability, utilization, and management of wild and semi-wild edible plants. The study area was stratified into two distinct agroecological zones, based on the framework by Hurni et al. (2016): the Lowland region (500-1500 meters) and the Midland region (1500-2300 meters). To ensure representative coverage, two districts from each zone were purposefully selected due to the known presence of wild and semi-wild edible plants. These districts were Tahtay Adiyabo and Tselemti in the Lowland region, and Tahtay Koraro and Medebay Zana in the Midland region (Figure 1). Within these districts, four peasant associations were randomly chosen: Adi Hageray (Tahtay Adiyabo), Mayteklit (Tselemti), Beles (Tahtay Koraro), and Bahra (Medebay Zana). The site selection was conducted in collaboration with researchers, woreda experts, development agents, and local administrators to ensure a diverse and representative sample.




Figure 1: Study area map

### 2.2. Sampling procedure and Sampling size determination

The study employed a mixed-methods approach to assess the availability, utilization, and management of wild and semi-wild edible plants across various community groups. A total of 20 key informants were purposefully selected, with the assistance of development agents,

from four study sites (5 informants per site). These key informants were instrumental in categorizing the wealth status of household participants, verifying recorded data on wild and semi-wild edible plants, and providing insights into the historical background of the community's utilization, conservation, and management practices related to these plants.

In addition, a random sample of 241 households was surveyed (as detailed in Table 1). The households represented a diverse cross-section of the community, including adults and elders, literate and illiterate individuals, women and men, as well as people from various socio-economic backgrounds (rich, medium wealth, and poor) and different religious affiliations. Sample size was calculated with the simple random sampling method based on proportional to household number presented below using formula (Kothari (2004)).

$$n = \frac{Z^2 * N * P * Q}{E^2 * (N - 1) + Z^2 * PQ} \dots \dots \dots Eq. (1)$$

Where, n=Sample size, N= Number of household, E= Allowed error which is 5 %=0.05, P=level of precision which is 5 %=0.05, Q=1-P=1-0.05=0.95, Z<sup>2</sup>=Confident interval 95, 1.96 from Z table

Table 1: lists of selected household in the study area

List of selected Kebele	Agroecology	No of Household	Sample size	Gender	
				Male	Female
Bahra	Midland	2097	59	54	5
Beles	Midland	2275	64	34	20
Adi Hageray	Lowland	1450	48	38	10
May Tekelit	Lowland	2488	70	48	22

**2.3. Data collection**

The study utilized checklists and semi structured interview questionnaires as data collection tools. Information was collected on various aspects of wild and semi-wild edible plant species, including their growth forms, utilized parts, natural habitats, propagation methods, fruit harvesting techniques, annual yields, conservation statuses, consumption patterns, market value, management practices and major threats to their survival. These key informations were collected from focus group discussion, key informant interview, household interview and filed observation.

**2.4. Data analysis**

The data were analyzed using descriptive statistics with Microsoft Excel and Stata software version 15. Closed-ended questions were examined using tables and percentages to enable result comparisons, while open-ended responses, including interviews and observations, were analyzed descriptively.

### 3. Result and Discussion

#### 3.1. Demographic and socio economic characteristics of the households

Most respondents were male-headed households and married. In the lowland agroecology, 55% of respondents were illiterate, unable to read or write, compared to 42.4% in the midland agroecology. The remaining respondents had completed either primary or secondary education (Table 2). Farming was the primary occupation in the study area, with 100% of households in lowland regions and 93.22% in midland regions relying on agricultural practices (Table 1). The residents had lived in the area for periods ranging from 1 to 81 years, with farming experience spanning 10 to 60 years. Similar studies in other regions have reported that the majority of communities rely on subsistence farming as their primary income source. (Musa et al., 2023; Regassa et al., 2015).

Table 2: Respondents information and demographic characteristics

Variables		Lowland Agro ecology		Midland Agro ecology	
		Frequency	Relative Frequency (%)	Frequency	Relative Frequency (%)
Gender	Male	86	72.88	98	79.67
	Female	32	27.12	25	20.33
	Total	118	100	123	100
Age	20-30	14	11.86	16	13.01
	31-40	41	34.75	23	18.70
	41-50	20	16.95	29	23.58
	50-60	25	21.19	35	28.46
	>60	18	15.25	20	16.26
Marital status	Single	10	8.47	10	9.76
	Married	88	74.58	96	78.05
	Divorced	14	11.86	10	6.50
	Widowed	6	5.08	8	5.69
Family size	0-3	18	15.25	21	17.07
	4-6	47	39.83	54	43.90
	>7	53	44.92	48	39.02
Education	Illiterate	52	44.07	38	30.89
	Primary	42	35.59	60	48.78
	Secondary	24	20.34	25	20.33
Occupation	Farming	118	100	110	89.43

	Farming and Trade	0	0	8	6.50
	Farming and Guard	0	0	2	1.63
	Government employee	0	0	3	2.44
Wealth status	Poor	68	57.63	57	46.34
	Medium	30	25.42	35	28.46
	Rich	20	16.95	31	25.20

### 3.2. Identified wild and semi wild edible plants and life forms

The study area is endowed with various wild and semi-wild edible (WSWE) plants that serve as both timber and non-timber forest products. A total of 32 WSWE plant species were identified across both agroecologies: 8 in the lowland, 11 in the midland and 13 in both. The number of WSWE plants recorded was higher in the midland agroecology compared to the lowland agroecology. This finding aligns with those reported by Woldemedhin et al. (2021) in the Ensaro district of the Amhara region. The growth habits of WSWE plants in the study area reflect a clear dominance of tree species, which make up 65.62% of the total vegetation, followed by tree/shrub species at 18.75%, and shrubs at 15.63% (Table 3). This dominance of trees is consistent with findings from other studies (Anbessa et al., 2024; Emire et al., 2022; Musa et al., 2023; Tariku and Eyayu, 2017; Demise, 2020). However, this result contrasts with studies by Woldemedhin et al. (2021), Anbessa (2016), Lulekal et al. (2011), Regassa et al. (2015), Biri et al. (2024), and Tahir et al. (2023), which reported a dominance of shrubs over other life forms of WSWE plants.

The families Malvaceae and Moraceae had the highest number of species, with four species each (12.5%). These were followed by Arecaceae, Ebenaceae, and Rhamnaceae, each with two species (6.25%). The remaining families were represented by only one species each. The dominance of species from the Moraceae family in the study area is consistent with findings from other parts of Ethiopia (Girmay et al., 2022; Regassa et al., 2015; Tariku and Eyayu, 2017). Additionally, the prevalence of Malvaceae and Moraceae, followed by Arecaceae, Rhamnaceae, Boraginaceae, Anacardiaceae and Rubiaceae, aligns with observations by Dejene et al. (2020). The number of WSWE plants identified in this study is comparable to those reported in other regions of Ethiopia (Musa et al., 2023), but it is lower than the numbers documented by Anbessa et al. (2024), Dejene et al. (2020), Biri et al. (2024), Tariku and Eyayu (2017), and Emire et al. (2022). The lower number of WSWE plants found in this study may be attributed to climatic and environmental conditions that could have limited the availability of wild edible plants, as well as differences in traditional knowledge and practices related to the use of WSWE plant species.

Table 3: List o of the WSWE plants and life form

Scientific name	Local name	Family	Edible	Life form
<i>Hyphaene thebaica</i>	Arkokobay	Arecaceae	Fruit	Tree
<i>Dovyalis abyssinica</i>	Ayahada	Salicaceae	Fruit	Tree/shrub
<i>Adansonia digitata L.</i>	Dima	Malvaceae	Fruit	Tree

	Gaba adgi		Fruit	Tree
<i>Grewia villosa</i>	Haben	Malvaceae	Fruit	Tree
<i>Tamarindus indica</i>	Humer	Fabaceae	Fruit	Tree
	Meshengoga		Fruit	Shrub
<i>Balanites aegyptica</i>	Meqie	<i>Zygophyllaceae</i>	Fruit	Tree
<i>Euclea schimperi</i>	Kiliow	Ebenaceae	Fruit	shrub
<i>Syzygium guineense</i>	Liham	Myrtaceae	Fruit	Tree
<i>Phoenix reclinata</i>	Siye	Arecaceae	Fruit	Tree
<i>Rhus natalensis</i>	Tetaelo	Anacardiaceae	Fruit	Tree/shrub
<i>Grewia ferruginea</i>	Tsimquya	Malvaceae	Fruit	Tree
<i>Vangueria edulis</i>	Guramayle	Rubiaceae	Fruit	shrub
<i>Ficus ingens/glutinosa</i>	Tsekente	Moraceae	Fruit	Tree
<i>Securinega virosa</i>	Harmazo	Phyllanthaceae	Fruit	Shrub
<i>Ficus sur</i>	Kodo	Moraceae	Fruit	Tree
	Wembley		Fruit	Tree
<i>Ziziphus abyssinica</i>	Abetere	Rhamnaceae	Fruit	Tree/shrub
<i>Carissa edulis</i>	Agam	Apocynaceae	Fruit	Tree/shrub
<i>Diospyros mespiliformis</i>	Aye	Ebenaceae	Fruit	Tree
<i>Ficus vasta</i>	Daero	Moraceae	Fruit	Tree
<i>Cordia africana</i>	Awhi	Boraginaceae	Fruit	Tree
<i>Ziziphus spina christi</i>	Gaba	Rhamnaceae	Fruit	Tree/shrub
<i>Grewia bicolor</i>	Mesoqa	Malvaceae	Fruit	Shrub
<i>Ximenia americana</i>	Milio	Olacaceae	Fruit	Tree
<i>Ficus sycomorus</i>	Sagla	Moraceae	Fruit	Tree
<i>Mimusops kummel</i>	Kumel	Sapotaceae	Fruit	Tree
<i>Moringa oleifera Lam</i>	Moringa	<i>Moringaceae</i>	Leaf	Tree
<i>Verbascum sinaiticum</i>	Tinkako	<i>Scrophulariaceae</i>	Fruit	Tree
	Harenker		Fruit	Tree/shrub
	Mulo kak		Fruit	Tree

### 3.3. Habitat, propagation and conservation status of WSWE

#### 3.3.1. Natural habitats

Natural forests, grasslands, riverine environments, protected areas, farmlands, shrublands, woodlands, and wetlands are home to various WSWE plants in Ethiopia. The results indicated that WSWE plants exist in different habitat types (Table 4). The majority of these species were found in riverine areas (27.95%), followed by woodlands (16.76%), exclosures (9.94%), home gardens (8.07%), and farmlands (8.07%) in the lowland area. In the midland area, the majority were found in riverine habitats (28%), followed by woodlands (13.71%), home gardens (13.15%), exclosures (9.2%), and farmlands (5.75%). Other habitats are indicated in Table 3. Previous studies have also identified farmlands as a low reservoir for WSWE plants,

whereas scrublands and woodlands were found to be the primary reservoirs for these plants (Woldemedhin et al., 2021).

Table 4: Natural habitats of the WSWE plants in the study area

Habitat	Lowland Agroecology		Midland Agroecology	
	Frequency	Relative frequency	Frequency	Relative frequency
Woodland		16.76		13.71
Farm land		8.07		5.75
Riverine		27.95		28.
Home garden		8.07		13.15
Exclosure		9.94		9.20
woodland and Farmland		1.86		2.99
woodland and riverine		1.86		5.75
Farm land and homestead		0.62		2.53
woodland and homestead		0.62		0.92
woodland, farmland, riverine and homestead		8.70		5.52
woodland, farmland and riverine		0.62		0.69
woodland, farm land and homestead		1.24		2.07
Homesteaded and exclosure		10.56		7.42
Riverine and homestead		1.24		0.69
Riverine and exclosure		1.86		1.61
Total		100		100

### 3.3.2. Propagation methods of WSWE

Proper propagation techniques are essential for maintaining plant populations, restoring degraded habitats, and ensuring a continuous supply of edible, medicinal, and culturally significant plants. In the study area, 100% of respondents in the lowland region and 96.88% in the midland region propagate WSWE plants using seeds (Table 5). In addition to seed propagation, *Ziziphus spina-christi* is propagated using rhizomes, while *Ficus sycomorus* and *Ficus vasta* are propagated through cuttings.

Table 5: Propagation methods of WSWE

Propagation methods	Lowland Agroecology		Midland Agroecology	
	Frequency	Relative frequency (%)	Frequency	Relative frequency (%)
Seed	20	100		96.88
Cuttings				0.46

Rhizome				1.84
Seed and cutting				0.23
Seed and rhizome				0.46
Seed and grafting				0.23

### 3.3.3. Conservation status and demands

WSWE plants are essential for biodiversity, food security, and medicinal and cultural values, and they require dedicated conservation measures. In the study area, however, the conservation status of WSWE plants faces mounting threats from habitat loss, overharvesting, climate change, agricultural expansion, population pressure, and charcoal production. A significant percentage of respondents 88.52% in the lowland and 78.30% in the midland agroecology, indicated an urgent need for conservation efforts for WSWE plants (Figure 2). Similar studies in other areas have also recommended implementing comprehensive conservation measures, as well as the domestication and cultivation of WSWE plants (Anbessa et al., 2024; Biri et al., 2024; Dejene et al., 2020; Musa et al., 2023; Tariku and Eyayu, 2017).




Figure 2: current status and conservation demands of the WSWE in lowland and midland agroecology

### 3.4. Harvesting Methods of the WSWE fruits

In the study area, the community employs various WSWE fruit harvesting techniques, including plucking, shaking the tree, cutting branches, using a long stick to pick the fruit, and throwing stones to dislodge the fruit. Among these, plucking, followed by picking the fruit with a long stick and shaking the tree, was the most common method in both agroecologies (Table 6). This finding aligns with the research of Yiblet and Adamu (2023) and Ojelel et al. (2019), who reported that plucking was the most common harvesting technique in Tach Gayint district, Amhara region, Ethiopia, and the Teso-Karamoja region, Uganda,

respectively. The use of techniques such as branch cutting, stoning, and using long sticks to pick fruit may hinder the growth and survival of the mother tree. Additionally, these methods can reduce fruit quality by damaging both ripe and unripe fruits, increasing their exposure to diseases and pathogens.

Table 6: Harvesting techniques of the WSWE

Fruit harvesting Techniques	Lowland Agroecology		Midland Agroecology	
	Frequency	Relative frequency	Frequency	Relative frequency
Plucking		67.40		59
Shaking		11.18		1.15
Picking by stick		11.80		10.12
From felled		0.62		3.45
Cutting branches		2.17		2.59
Plucking and shaking		1.86		2.53
Plucking and picking by stick		1.24		8.97
Plucking and from felled		0		0.69
Plucking, picking by stick and shaking		0		9.2
Plucking, shaking and from felled		0		0.92
Picking by stick and from felled		0		0.69
Plucking, shaking, picking by stick and from felled		0		0.23
Plucking , picking by stick and from felled		0		0.23
Shaking and from felled		0		0.23
Shaking and picking by stick		3.11		0
Stoning		0.62		0

### 3.5. Fruit collector and seller groups

In the study area, the results showed that household members participated in WSWE fruit collection, with children being the primary fruit collectors, followed by other household members (Table 7). Consistent with these findings, women and children (or youngsters) were documented as the major WSWE fruit gatherers, followed by other household members (Biri et al., 2024; Musa et al., 2023; Regassa et al., 2015). Demise (2020) also reported that younger age groups were more actively involved in wild edible fruit collection compared to other household members. Additionally, the market assessment indicated that children were the primary sellers of WSWE fruits in the local markets of the study area (Table 6). The low participation of household heads in selling fruits may be attributed to cultural influences, community perceptions, and individual interests.

Table 7: WSWE fruit collector and seller groups

	Household members	Lowland Agroecology		Midland Agroecology	
		Frequency	Relative frequency (%)	Frequency	Relative frequency (%)
Fruit collectors	Husband	0	0		0.46
	Children		85.1		55.24
	Women and children's		2.48		7.59
	Husband, Women and children		12.42		36.09
	Husband and children's	0	0		0.62
Fruit sellers	Husband	0	0		0.46
	Women		4.35		0.62
	Children		95.03		93.56
	Husband, Women and children		0.62		5.13
	Husband and Women	0	0		0.23

### 3.6. Plant parts used and consumption mode

Fruits, seeds, and leaves are the edible parts of WSWE plants recorded in the study area. Fruits were the most commonly consumed part, constituting 96.6% of consumption, followed by leaves at 3.1% (Figure 3). All household members consumed WSWE fruits during times of food shortage and in normal seasons. In contrast to our study area, people in other parts of Ethiopia also utilize the roots and barks of WSWE plants, in addition to the fruits, seeds, and leaves (Musa et al., 2023). This suggests that different regions have distinct cultural practices and traditional knowledge regarding the use of WSWE plants. Overall, many studies have shown that fruit is the major part consumed by communities (Anbessa, 2016; Anbessa et al., 2024; Musa et al., 2023; Regassa et al., 2015; Guzo et al., 2023; Emire et al., 2022; Demise, 2020), which is consistent with the present findings.




Figure 3: WSWE parts used for food consumption

The results on consumption modes showed that communities consumed WSWE fruits in various forms, including fresh ripe fruits, fruit juice, cooked/roasted fruits, cooked leaves, and raw seeds. Approximately 74.53% of the fruits in lowland areas and 97.01% in midland areas were consumed raw, without processing (Table 8). In addition to being eaten raw, *Tamarindus indica* was also consumed in cooked form, often combined with other ingredients, and as juice. *Mimusops kummel* was typically consumed in roasted form, while *Moringa oleifera* leaves were cooked, and *Cordia africana* seeds were eaten for medicinal purposes. Similar to the present findings, previous studies have also reported that a higher proportion of fruits were consumed raw (Anbessa et al., 2024; Demise, 2020; Dejene et al., 2020; Girmay et al., 2022; Musa et al., 2023; Tariku and Eyayu, 2017).

Table 8: Consumption mode of the WSWE fruits

Consumption mode	Lowland Agroecology		Consumption mode	Midland Agroecology	
	Frequency	Relative Frequency		Frequency	Relative Frequency
Fresh ripe fruit /Raw		74.53	Raw		97.01
Juice		16.15	Cooked/roasted		0.46
Dried		0.62	Dried		0.46
Raw and cooked		1.24	Raw and dried		2.07
Raw and dried		0.62			
Cooked and juice		5.59			
Raw, cooked and juice		0.62			
Seed		0.62	Seed		0
Total		100			100

### 3.7. Yield and market assessment of the commonly used WSWE

The seasonal yield of WSWE fruits per stem varied from plant to plant. The average yield ranged from 3 to 97.9 kilograms per stem, with the highest yield coming from *Syzygium guineense* and the lowest from *Verbascum sinaiticum* (Table 9). The market assessment revealed that, except for *Tamarindus indica*, *Balanites aegyptiaca*, *Ziziphus spina-christi*, *Syzygium guineense*, *Vangueria edulis*, *Diospyros mespiliformis*, *Cordia africana*, *Ximenia americana*, and *Mimusops kummel*, 75% of the WSWE fruits were not available for sale in the local markets (Table 8). Among these, *Tamarindus indica*, *Balanites aegyptiaca*, *Vangueria edulis*, *Hyphaene thebaica*, and *Adansonia digitata* were sold individually by count, while *Diospyros mespiliformis*, *Ziziphus spina-christi*, *Cordia africana*, *Ximenia americana*, and *Mimusops kummel* were typically sold using traditional local measuring units, such as "Tanika" (where 2 Tanika equal 1 kilogram). Similar to this finding, previous studies conducted in other regions have shown that a higher number of WSWE fruit trees are not sold in markets as a food source (Alemayehu et al., 2015; Anbessa, 2016; Emire et al., 2022; Kidane et al., 2014). In contrast, research carried out in the neighboring country of Sudan revealed that WSWE fruits contribute 50-60% and 75-100% of the total annual family income in lowland and midland areas, respectively (Salih and Ali, 2014).

Table 9: Seasonal fruit yield of the WSWE/stem

Scientific name	Local name	Annual Yield (Kg)	Price/Kg or per fruit
	Abetere	29.9	0

	Agam	33.6	0
<i>Diospyros mespiliformis</i>	Aye	67.6	20-30 birr/Kg
<i>Cordia africana</i>	Awhi	77.7	20-35 Kg
	Daero	66	0
<i>Ziziphus spina-christi</i>	Gaba	59.8	25-40 Kg
	Kiliow	22.1	0
<i>Mimusops kummel</i>	Kumel	35.2	15-30 Kg
<i>Syzygium guineense</i>	Liham	97.9	20-30 Kg
	Milio	29.9	15-30 Kg
	Sagla	68.5	0
	Siye	11.3	0
	Tetaelo	12.08	0
	Tsinkuya	24.5	0
<i>Hyphaene thebaica</i>	Arkokobay	72.5	The price for one fruit is 5 birr
	Ayahada	10.8	0
	Dima	64.9	0
	Gaba adg	80	0
<i>Vangueria edulis</i>	Guramayl	15	The price for five fruit is 2 birr
	Haben	5	0
	Harenker	30.3	0
	Harmazo	20.9	0
<i>Tamarindus indica</i>	Humer	66.6	The price for five fruit is 1 birr
	Kiliow	22.1	0
	Kodo	55	0
<i>Balanites aegyptiaca</i>	Mekie	47.5	The price for five fruit is 2 birr
	Meshengo	7.1	0
	Mesoka	6.3	0
	Mulo kak	30	0
<i>Verbascum sinaiticum</i>	Tinkako	3	0
	Tsekente	5.75	0
	Wemblay	10	0
	Moringa		

### 3.8. Other multi-purpose values of WSWE Plants

In addition to supplementing the regular food supply and filling the food gap during times of famine or drought, WSWE plants in the study area serve multiple purposes. These include providing forage/fodder, medicine, fuel wood, and materials for agricultural tools, furniture, construction, and rope. WSWE plants are also used for shade, erosion control, fencing, and saddle making. Furthermore, they play a role in agroforestry, income generation, and are integral to religious and cultural ceremonies.

### 3.9. Side Effects of the WSWE

The results indicate that the majority of WSWE plants, specifically 96.27% in lowland areas and 92.87% in midland areas, do not have disease-causing effects on human health (Table 10). However, some WSWE plants can have negative effects when consumed in excessive amounts. For example, overconsumption of *Balanites aegyptiaca* may lead to diarrhea, *Diospyros mespiliformis* can cause constipation, urinary incontinence, and abdominal bloating, *Ximenia americana* may result in dehydration, heart disease, and itchy teeth, *Tamarindus indica* can cause itchy teeth and constipation, *Ziziphus spina-christi* may lead to abdominal bloating, and *Cordia africana* can cause nausea. Excessive intake or overdose of WSWE fruits can lead to side effects on human health (Anbessa, 2016).

Table 9: Overconsumption side effects of WSWE plants

Table 10: Side effects of The WSWE plants

Side effect	Lowland Agroecology		Side effect	Midland Agroecology	
	Frequency	Relative Frequency		Frequency	Relative Frequency
No side effect		96.27	No side effect		92.87
Abdominal bloating		0.62	Abdominal bloating		0.92
Itchy teeth		0.62	Itchy teeth		0.69
Diarrhea		1.87	Urinary incontinence		0.23
Constipation		0.62	Constipation		1.84
			Heart disease		0.23
			Abdomen pain		2.99
			Nausea		0.23
Total		100			100

### 3.9. Major threatening factors and management practices of the WSWE

#### 3.9.1. Major threatening factors

In the study area, people utilize a variety of WSWE plants from natural forests, homesteads, riverine areas, exclosures, and farmlands for purposes such as medicinal use, income generation, and food. However, despite their importance, the abundance and diversity of WSWE plants are decreasing due to anthropogenic and environmental threats. Respondents in the study area identified the main threats as agricultural expansion, charcoal production, and overgrazing. Specifically, in the lowland agroecology, 79.63% of respondents highlighted agricultural expansion and charcoal production, while 20.37% cited grazing as a major threat. In the midland agroecology, 76.32% pointed to agricultural expansion and charcoal production, and 23.68% identified grazing as a significant threat. These and other factors may contribute to the decline in the number of WSWE plants in the study area. Similar to these findings, many studies in other regions have identified agricultural expansion, overgrazing, and fuelwood production as the primary threats to WSWE plants (Kebebew and Mohamed, 2017; Girmay et al., 2022; Musa et al., 2023; Regassa et al., 2015; Tariku and Eyayu, 2017; Emire et al., 2022).

#### 3.9.2. Management and Conservation Practices

Effective management of WSWE plants includes various strategies, such as in situ conservation, selective thinning and pruning, habitat restoration through plantations and exclosures, community-driven harvesting guidelines and integration with agricultural systems.

Despite the numerous roles of WSWE plants in enhancing food security through consumption and income generation, providing stable ecological services, creating job opportunities and supporting religious and cultural practices in the study area, these economically important species have received little attention. Fifty and more than fifty percent of the respondents stated that there is improper WSWE plants utilization and no effective management practices have been implemented to conserve and restore these plants (Figure 4). Consequently, it becomes low diverse in the study area. Similar to the present result, previous studies conducted in other region also showed that little attention plus misuse causes to reduction of the WSWE plants (Anbessa, 2016; Guzo et al., 2023; Woldemedhin et al., 2021).




Figure 4: Management practices of the WSWE plants in the study area

#### 4. Conclusion

The study revealed that the northwestern Tigray region is endowed with a diverse range of WSWE plants, with a total of 32 species documented across both lowland and midland areas. The majority (96.88%) of these WSWE plants are native, with only a small proportion of exotic species (3.12%). Despite the abundance of WSWE plants, only a few are marketed as food sources. The primary edible parts consumed are fruits, which serve to supplement the local diet. However, excessive consumption of WSWE fruits has been linked to various health issues, including abdominal pain, bloating, heart disease, nausea, urinary discomfort, constipation, and itchy teeth.

Furthermore, local perceptions of WSWE plants are generally low, particularly in terms of management practices, parts consumed, sustainable utilization, and conservation strategies. Currently, the diversity and population of WSWE plants face threats from both anthropogenic and natural factors, such as agricultural expansion, illegal cutting for firewood and charcoal production, overgrazing, urbanization, and climate change. These factors are contributing to the declining availability of WSWE plants, potentially leading to their disappearance. Therefore, this study emphasizes the need to raise awareness among local communities about the sustainable and wise use of WSWE plants. It also calls for urgent restoration and conservation efforts to preserve these valuable resources for future generations.

## References

1. Abebe D and Ayehu A (1993) Medicinal Plants and Enigmatic Health Practices of Northern Ethiopia. Monograph, Addis Ababa,
2. Alemayehu, G., Asfaw, Z., and Kelbessa, E. (2015). Plant diversity and ethnobotany in Berehet District, North Shewa Zone of Amhara Region (Ethiopia) with emphasis on wild edible plants. *Journal of Medicinal Plants Studies*, 3(6), 93-105. *Journal of Medicinal Plants Research* 3 (6): 93-105.
3. Amente, D. A. (2017). Ethnobotanical survey of wild edible plants and their contribution for food security used by Gumuz people in Kamash Woreda; Benishangul Gumuz Regional State; Ethiopia. *Journal of Food and Nutrition Sciences*, 12(1), 217-224. DOI: 10.11648/j.jfns.20170506.12
4. Anbessa, B. (2016). Ethnobotanical study of wild edible plants in Bule Hora Woreda, Southern Ethiopia. *African Journal of Basic & Applied Sciences*, 8(4), 198-207. ISSN 2079-2034, DOI: 10.5829/idosi.ajbas.2016.8.4.23529
5. Anbessa, B., Lulekal, E., Getachew, P., and Hymete, A. (2024). Ethnobotanical study of wild edible plants in Dibatie district, Metekel zone, Benishangul Gumuz Regional State, western Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 20(1), *Journal of Ethnobiology and Ethnomedicine* <https://doi.org/10.1186/s13002-024-00671-2>
6. Asfaw, Z. (2008, March). The future of wild food plants in southern Ethiopia: ecosystem conservation coupled with enhancement of the roles of key social groups. In *International Symposium on Underutilized Plants for Food Security, Nutrition, Income and Sustainable Development* 806 (pp. 701-708). DOI: [10.17660/ActaHortic.2009.806.87](https://doi.org/10.17660/ActaHortic.2009.806.87)
7. Belete, G., Asfaw, Z., and Tamirat, T. (2021). Diversity of wild edible fruit bearing woody species in different land use and management system in Dangur district, north western Ethiopia. *Journal of Global Ecology and Environment*, 12-22. SSN: 2454-2644
8. Berihun, T., and Molla, E. (2017). Study on the diversity and use of wild edible plants in Bullen istrict Northwest Ethiopia. *Journal of Botany*, 2017(1), 8383468. <https://doi.org/10.1155/2017/8383468>
9. Biri, S., Ayenew, B., Dida, G., Sebsibe, A., Gurmessa, F., Woldeab, B., ... and Megersa, M. (2024). Ethnobotanical study of wild edible plants in Arsi Robe district of East Arsi Zone, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 20(1), 70. <https://doi.org/10.1186/s13002-024-00703-x>
10. Dejene, T., Agamy, M. S., Agúndez, D., and Martin-Pinto, P. (2020). Ethnobotanical survey of wild edible fruit tree species in lowland areas of Ethiopia. *Forests*, 11(2), 177, DOI: 10.3390/f11020177
11. Demise, S. (2020). Ethno botanical study of wild edible plants in Adola district, southern, Ethiopia. *IJRAR-International Journal of Research and Analytical Reviews (IJRAR)*, 7(2), 212-228., E-ISSN 2348-1269
12. Emire, A., Demise, S., Giri, T., and Tadele, W. (2022). Ethnobotanical study of wild edible plants in Liben and Wadera Districts of Guji Zone, Southern Ethiopia. *Glob J Agric Res*, 10(3), 47-65. Print ISSN: 2053-5805

13. Giday, M., and Teklehaymanot, T. (2023). Use of wild edible and nutraceutical plants in Raya-Azebo District of Tigray Region, northern Ethiopia. *Tropical Medicine and Health*, 51(1), 58. <https://doi.org/10.1186/s41182-023-00550-8>
14. Girmay, M. G., Lulekal, E., Belay, B., and Gebrehiwot, K. (2022). Wild edible plants study in a dryland ecosystem of Ethiopia. *Daagu International Journal of Basic and Applied research (DIJBAR)*, 4(2), 105-119.
15. Guinand, Y., and Lemessa, D. (2001). Wild-food plants in Ethiopia: Reflections on the role of wild foods and famine foods at a time of drought. *The potential of indigenous wild foods*, 22, 31.
16. Guzo, S., Lulekal, E., and Nemomissa, S. (2023). Ethnobotanical study of underutilized wild edible plants and threats to their long-term existence in Midakegn District, West Shewa Zone, Central Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 19(1), 30. <https://doi.org/10.1186/s13002-023-00601-8>
17. Kothari CR (2004) Research methodology: Methods and techniques. New Age International Publishers, New Delhi, India.
18. Kebebew, M., and Mohamed, E. (2017). Indigenous knowledge on use of medicinal plants by indigenous people of Lemo district, Hadiya zone, Southern Ethiopia. *International Journal of Herbal Medicine*, 5(4), 124-135. E-ISSN: 2321-2187, P-ISSN: 2394-0514
19. Kidane, B., Van der Maesen, L. J. G., Van Andel, T., Asfaw, Z., and Sosef, M. S. M. (2014). Ethnobotany of wild and semi-wild edible fruit species used by Maale and Ari ethnic communities in southern Ethiopia. *Ethnobotany Research and applications* 12:455-471
20. Lulekal, E., Asfaw, Z., Kelbessa, E., and Van Damme, P. (2011). Wild edible plants in Ethiopia: a review on their potential to combat food insecurity. *Afrika focus*, 24(2), 71-122.
21. Musa Abdella, Bira Cheneke and Megarsa Ketama (2023). Assessment of Wild Edible Plant Species in East Hararghe Zone, Oromia, Ethiopia. *International Journal of Science, Technology and Society* 2023; 11(3): 81-94, DOI: 10.11648/j.ijsts.20231103.11
22. Ojelel, S., Mucunguzi, P., Katuura, E., Kakudidi, E. K., Namaganda, M., and Kalema, J. (2019). Wild edible plants used by communities in and around selected forest reserves of Teso-Karamoja region, Uganda. *Journal of ethnobiology and ethnomedicine*, 15, 1-14. <https://doi.org/10.1186/s13002-018-0278-8>
23. Regassa, T., Kelbessa, E., and Asfaw, Z. (2015). Ethnobotany of wild and semi-wild edible plants of Chelia District, West-Central Ethiopia. *Science, Technology and Arts Research Journal*, 3(4), 122-134. DOI: [10.4314/star.v3i4.18](https://doi.org/10.4314/star.v3i4.18)
24. Ruffo, C. K., Birnie, A., and Tengnäs, B. (2002). Edible wild plants of Tanzania. Published by the Regional Land Management Unit, RELMA/Sida, ICRAF House, Gigiri, Nairobi, Kenya
25. Salih, N. K. E. M., and Ali, A. H. (2014). Wild food trees in Eastern Nuba Mountains, Sudan: Use diversity and threatening factors, *Journal of Agriculture and Rural Development in the Tropics and Subtropics*, Vol. 115 (1), 1–7, urn:nbn:de:hebis:34-2014020344903

26. Seyoum, Y., Teketay, D., Shumi, G., and Wodafirash, M. (2015). Edible wild fruit trees and shrubs and their socioeconomic significance in central Ethiopia. *Ethnobotany Research and Applications*, 14, 183-197. <http://dx.doi.org/10.17348/era.14.0.183-197>
27. Tahir, M., Abraham, A., Beyene, T., Dinsa, G., Guluma, T., Alemneh, Y and Mohammed, A. (2023). The traditional use of wild edible plants in pastoral and agro-pastoral communities of Mieso District, eastern Ethiopia. *Tropical Medicine and Health*, 51(1), 10. <https://doi.org/10.1186/s41182-023-00505-z>
28. Tariku Berihun and Eyayu Molla (2017). Study on the Diversity and Use of Wild Edible Plants in Bullen District, Northwest Ethiopia, *Hindawi Journal of Botany*, <https://doi.org/10.1155/2017/8383468>
29. Teklehaymanot, T., and Giday, M. (2010). Ethnobotanical study of wild edible plants of Kara and Kwego semi-pastoralist people in Lower Omo River Valley, Debub Omo Zone, SNNPR, Ethiopia. *Journal of ethnobiology and ethnomedicine*, 6(1), 1-8., <http://www.ethnobiomed.com/content/6/1/23>
30. Woldemedhin, A. A., Lulekal, E., Bekele, T., Debella, A., Debebe, E., and Tessema, S. (2021). Ethnobotanical study of edible wild plants in Ensaro district, Amhara regional state, Ethiopia. Research square, DOI: [https://doi.org/10.21203/rs-418877/v1](https://doi.org/10.21203/rs.3.rs-418877/v1)
31. Yiblet, Y., and Adamu, E. (2023). An ethnobotanical study of wild edible plants in Tach Gayint district, South Gondar zone, Amhara region, Northwestern Ethiopia. *Evidence-Based Complementary and Alternative Medicine*, (1), 7837615. <https://doi.org/10.1155/2023/7837615>