

Impact of Organic inputs (Jeevamruth and Bijamruth) on soil nutrients and uptake by Fenugreek

Abstract:

The present study on 'Biochemical evaluation of organic inputs (Jeevamruth and Beejamruth) and their efficacy on Greens' was carried out at the Department of sustainable organic agriculture, Tamil Nadu Agricultural University, Coimbatore and the laboratory experiments were carried out at the Department of Environmental Science. A field experiment was conducted in a randomized block design with three replications. Biometric observations were taken during ~~the~~ 10th, 20th and 30th days after sowing. Soil samples were collected during ~~the~~ initial and at the stage of harvest and analysed for chemical and biological properties. Plant samples were analysed for protein and chlorophyll contents. The application of Jeevamruth in soil resulted in an increase in the nutrients (NPK) and organic carbon values. Liquid organic preparations contain ~~a~~ higher number of bacteria, fungi, and actinomycetes. From the studies, it is evident that Beejamruth is to be used on the day of preparation while Jeevamruth should be used within 10 to 15 days from the date of preparation. The application of these liquid formulations would supplement the nutrients when biofertilizers were applied.

Key-words: Jeevamruth, Beejamruth, Biochemical evaluation, Organic inputs, Organic farming, Green revolution

Introduction:

~~Green~~ ~~The green~~ revolution ~~had~~ ~~has~~ intensified agriculture to meet the ~~ever~~ ~~ever~~ increasing demand for food and fibre, which is a practice at great cost to the environment resulting in continuous loss of natural ecosystems, ground-water, food-stuff pollution and other environmental degradation. The greatest challenge to be faced by the nation in the coming years is, to provide safe food for the growing population in the country. In this regard, organic farming which is a holistic production management system for promoting and enhancing ~~the~~

health of ~~agro-agro~~-ecosystem, has gained wide recognition as a valid alternative to conventional food production and ensures safe food for human consumption. This farming system avoids large use of synthetic fertilizers, growth regulators, livestock feed additives and relies on green manures, crop rotations, crop residues, animal manures, biofertilizers, different kinds of ~~cow-cow~~-based liquid organic manures such as Panchagavya, Jeevamruth, Beejamruth, Amritpani etc. Among the liquid formulations, panchagavya is one of the most important liquid formulations to be considered for the study of shelf life as it not only proves to be most effective for crop growth but is also widely used by ~~the~~ farmers (Sugumaran *et al.*, 2018). Compounds Erioflorin and nagilactone A present in Beejamruth are plant growth regulators and hence Beejamruth is a valid and effective alternative fertilizer for the production of safe and good quality food ~~produces-products~~ which could cater to the needs of ~~the~~-modern Indian requirements (Goveanthan *et al.*, 2019). Compounds Isoenanthic acid, Columbianetin and Lomatin, 1, 6-Hexanediol, Mevastatin and Gitoxigenin, Dibutoxy anthracine, Erioflorin and nagilactone, Trimegestonea, Rofe Coxib, Clupanodonic acid are present in Jeevamruth which helps in the plant metabolism and also improves its growth (Goveanthan *et al.*, 2021).

Organic agriculture is now finding a place in the mainstream of development and shows great promise commercially, socially and environmentally. While there is a continuum of thought from earlier days to the present, the modern organic movement is radically different from its original form. Liquid formulations that are used in organic agriculture like Panchagavya, Beejamruth and Jeevamruth are ~~the~~ fermented products ~~which-that~~ are used as plant ~~growth-growth~~-enhancing substances prepared with material available ~~with-to~~ farmers. They are ~~the~~-rich sources of beneficial micro-flora ~~which-that~~ supports, and stimulates ~~the~~-plant growth and helps in getting better vegetative growth and also good quality yield. Formulations prepared from agricultural by-products *viz.*, bran of grains, oil cakes, farmyard manure etc., ~~which~~-are found to support excellent growth carrier and storage media (Devakumar *et al.*, 2011). During the last few years, there has been an increasing interest in the use of Panchagavya, Beejamruth, Jeevamruth and other liquid organic formulations in organic agriculture.

Materials and methods:

Field experimental details

The experiment was conducted in a randomised block design with three replications. The experimental layout was kept undisturbed throughout the period of investigation and fenugreek seeds were soaked for 1 hour in the Beejamruth solution before sowing and were sown in the field.

Design : RBD

Number of treatments : 7

Number of replications : 3

The treatment details are given below:

Treatments

T₁–Control

T₂-Jeevamruth @ 3% Spray (Green gram flour)

T₃-Jeevamruth @ 5 % Spray (Green gram flour)

T₄-Jeevamruth @ 3 % Spray (Black gram flour)

T₅-Jeevamruth @ 5 % Spray (Black gram flour)

T₆-Jeevamruth @ 3 % Spray (Green gram + Black gram flour)

T₇-Jeevamruth @ 5 % Spray (Green gram +Black gram flour)

Preparation of soil sample

Pre-experiment composite soil samples were collected and analysed for mechanical and chemical properties. Post-harvest soil samples were collected from each treatment plot after the harvest of Fenugreek and analysed for the pH, EC, organic carbon and major nutrients by following the standard procedures.

Preparation of the plant sample

The samples were collected for the estimation of dry matter production used for the estimation of nutrient uptake. The ~~oven-oven-~~dried plant samples were ground using Wiley-Mill, sieved and analysed to calculate the total NPK up taken by multiplying the contents of N, P and K with the dry matter at the respective stages, the uptake was arrived at and reported. The samples were pulverized and sieved through a 0.2 mm mesh sieve and the analysis ~~were-was~~ carried out by following standard procedures.

Results and Discussion:

The results from the experiment on “Biochemical evaluation of organic inputs (Jeevamruth and Beejamruth) and their efficacy on greens “conducted at the Department of Sustainable Organic Farming, Tamil Nadu Agricultural University, Coimbatore are summarized below.

Soil parameters

The soil pH was not significantly influenced by the different treatments. However, numerically higher soil pH was observed as given in ~~the~~ (Table 1). The Jeevamruth, as an organic source of nutrients slightly increased the soil pH. ~~But-However,~~ no significant difference in pH among various treatments was noticed. However, the treatment ~~which-that~~ received Jeevamruth @ 5 % Spray (Green gram flour) recorded higher pH among the treatments. This was supported by Elias Azar (1980) who reported that soil pH had increased due to the application of poultry manure as an organic nutrient source. The soil EC was not significantly influenced by the different treatments. The high-~~low~~ EC value was found in Jeevamruth @ 5 % Spray (Green gram flour) and Jeevamruth @ 5 % Spray (Black gram flour). At the time of the inception of the study, the soil organic content was 0.28 per-cent. After harvest of the fenugreek crop, the organic carbon content of the soil was higher in Jeevamruth @ 5% Spray (Green gram flour) (1.08%), followed by Jeevamruth @ 5 % Spray (Green gram +Black gram flour) (0.92%). The lowest organic carbon content was observed in the control (0.33%). ~~Addition-The addition~~ of Jeevamruth @ 5 % Spray (Green gram flour) resulted in a significant increase in the organic carbon content of the soil-. These results are in agreements with the findings of Mathan (2000) and Maskina *et al.* (1988).

The available soil nitrogen was not significantly influenced by the sources of nutrition. Numerically higher soil nitrogen was recorded in the treatments Jeevamruth @ 3% Spray (Green gram flour) and Jeevamruth @ 5 % Spray(Green gram +Black gram flour). The highest soil available phosphorus at 30 DAS, was recorded in the Jeevamruth @ 5% Spray (Green gram flour) and Jeevamruth @ 5 % Spray(Green gram +Black gram flour) treatments with values of 19.0 and 16.0 kg ha⁻¹ respectively. The high soil potassium content was recorded in Jeevamruth @ 5% Spray (Green gram flour) (264.0 kg ha⁻¹) followed by Jeevamruth @ 5 % Spray(Green gram +Black gram flour) (239.0 kg ha⁻¹) treatments.

Plant Total NPK

The analysis for total NPK in fenugreek after harvest showed that the total nitrogen content was found to be high in Jeevamruth @ 5 % Spray(Green gram +Black gram flour) with 1.5 per-cent, whereas the total phosphorus content was high in Jeevamruth @ 5 % Spray(Green gram) (0.33 per-cent). The potassium content was high in the treatments Jeevamruth @ 5 % Spray(Green gram), Jeevamruth @ 5 % Spray (Black gram flour), and Jeevamruth @ 5 % Spray(Green gram +Black gram flour) with 0.22 per-cent and they were on par with each other (Table 3). Nutrient uptake is a cordially event of nutrient concentration and dry matter accumulation. Organic manures promoted nutrient utilization and ~~accounts~~ accounted for better NPK uptake. Increased uptake might be due to the higher availability of nutrients from the soil reservoir and also from the added sources of organic manures (Priyadarsini and Prasad, 2003). The rate of uptake is dependent upon ~~by~~ crop N demand, phonological stage, soil N availability, transpiration, rooting depth and soil water status. Crop nitrogen demand is estimated depending on the rate of growth and the maximum concentration of nitrogen that different organic can accumulate depends upon their composition. Nitrogen uptake was maximum in Jeevamruth @ 5 % Spray(Green gram) and it was comparable with Jeevamruth @ 5 % Spray(Green gram +Black gram flour). Organic inputs are known to have a favorable effect on soil structure, texture and tilth thus ~~facilitate~~ facilitating quick and greater availability of plant nutrients and ~~provides~~ providing a better environment for root growth and proliferation, thereby creating a more absorptive surface for uptake of nutrients. These results ~~are in conformity~~ conform with the findings of Chavan *et al.* (1997); Shashidhara (2000) and Kuttimani (2004) in chilies. The organic inputs might have increased the soil organic P content leading to increased P availability. Higher phosphorus uptake was recorded in T₃ which was on par with T₇. Increased P availability might be due to the ~~solubilisation~~ solubilization of

native P by the organic acids produced during organic inputs decomposition, thus leading to better utilization of available P, which in turn favored better P uptake. Similar results were also obtained by Beulah (2001), Sreekhantan (1987) and Somasundaram (1991). The highest K uptake was registered with Jeevamruth @ 5 % spray (Use Green gram flour for preparation). The lower level of K uptake was observed in control (no manure/ no spray) at all the growth stages of the crop. The increased uptake of K observed in above said treatments might be the result of increased availability of K in soils due to the basal application of enriched farm yard manure. The enhanced K availability irrespective of the season coupled with higher K uptake due to organic manure incorporation could be attributed to higher DMP and K absorption, evidencing the priming effect of K contribution by organic manure. These results ~~are in conformity~~ conform with the findings of Santos *et al.*(1990) and Kuttimani (2004). The plant height, root length, and single plant weight in fenugreek are high in the treatment as Jeevamruth 5% spray was observed as a viable organic approach to improve soil and eco-friendly fenugreek production (Goveanthan *et al.*, 2020). In another study, seeds treated with Panchagavya and Jeevamruth separately, the maximum shoot length and root length were recorded in Panchagavya treatment, and minimum shoot length and root length ~~of was~~ recorded in Jeevamruth treated seeds and also the panchagavya treated seeds registered the maximum vigor index (Akila *et al.*, 2020). In another study, the soluble protein content (0.87 mg/g) and total sugar content (11.20 µg/g) were found to be improved in Panchagavya (Groundnut cake instead of ghee) @ 3% spray treatment and Recommended dose of NPK fertilizer applied plants (Sugumaran *et al.*, 2019).

Table 1. Effect of organic nutrient source (Jeevamruth) on soil pH, EC (dS m⁻¹) and organic carbon (%)

Treatment	pH	EC (dSm⁻¹)	Organic Carbon (%)	
T ₁ - Control	7.26	0.3	0.33	
T ₂ - Jeevamruth @ 3% Spray(Green gram)	7.28	0.8	1.08	
T ₃ - Jeevamruth @ 5 % Spray(Green gram)	7.29	1.5	0.90	
T ₄ - Jeevamruth @ 3 % Spray(Black gram)	7.40	0.4	0.46	
T ₅ - Jeevamruth @ 5 % Spray(Black gram)	7.62	1.5	0.72	
T ₆ - Jeevamruth @ 3 % Spray(Green gram + Black gram)	7.58	0.4	0.52	
T ₇ - Jeevamruth @ 5 % Spray(Green gram + Black gram)	7.74	0.8	0.92	
	SEd	0.0021	0.0055	0.003
	CD (P = 0.05)	0.0045	0.0119	0.06

Table 2. Effect of organic nutrient source (Jeevamruth) on soil available nitrogen, available phosphorus, available potassium (kg ha-1)

Treatment	Nitrogen (Kg ha⁻¹)	Phosphorus (Kg ha⁻¹)	Potassium (kg ha⁻¹)
T ₁ - Control	248.0	7.0	122.0
T ₂ - Jeevamruth @ 3% Spray(Green gram)	284	11	232
T ₃ - Jeevamruth @ 5 % Spray(Green gram)	392	19	264
T ₄ - Jeevamruth @ 3 % Spray(Black gram)	300	10	182
T ₅ - Jeevamruth @ 5 % Spray(Black gram)	348	15	198
T ₆ - Jeevamruth @ 3 % Spray(Green gram + Black gram)	308	12	188
T ₇ - Jeevamruth @ 5 % Spray(Green gram + Black gram)	360	16	239
SEd	0.5304	0.0438	0.5039
CD(P = 0.05)	1.1556	0.0954	1.0980

Table 3. Effect of organic nutrient source (Jeevamruth) on Fenugreek Plant Total nitrogen, Total phosphorus, Total potassium (%)

Treatment	Nitrogen (%)	Phosphorus (%)	Potassium (%)
T ₁ - Control	0.16	0.05	0.17
T ₂ - Jeevamruth @ 3% Spray(Green gram)	0.70	0.12	0.20
T ₃ - Jeevamruth @ 5 % Spray(Green gram)	1.23	0.33	0.23
T ₄ - Jeevamruth @ 3 % Spray(Black gram)	1.00	0.16	0.20
T ₅ - Jeevamruth @ 5 % Spray(Black gram)	1.50	0.25	0.22
T ₆ - Jeevamruth @ 3 % Spray(Green gram + Black gram)	1.06	0.18	0.21
T ₇ - Jeevamruth @ 5 % Spray(Green gram + Black gram)	1.50	0.31	0.22
SEd	0.0051	0.0011	0.0002
CD(P = 0.05)	0.0112	0.0024	0.0005

Conclusions:

From the field experiment can be ~~is~~ inferred that ~~the~~ the plant height, root length and single plant weight are highest in the treatment T₃(Jeevamruth @ 5% spray). The application of Jeevamruth in soil resulted in an increase in the nutrients (NPK) and organic carbon values. Liquid organic preparations contain ~~a~~ higher number of bacteria, fungi, and actinomycetes. From the studies ~~it~~ it is evident that Beejamruth is to be used on the day of preparation while Jeevamruth should be used within 10 to 15 days from the date of preparation. The application of these liquid formulations would supplement the nutrients when biofertilizers were applied.



Fig 1. Experimental field view

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests ~~OR~~ non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

References:

- Akila, S., Sugumaran, M. P., Suganya, K. and Somasundaram. E. (2020). Studies on Testing the Efficacy of Liquid Organic Inputs (Panchagavya and Jeevamruth) on

- Maize (*Zea Mays* L.) Germination. Current Journal of Applied Science and Technology, 39 (23),134-37.
- Beulah, A. 2001. Growth and development of moringa (*Moringa oleifera* Lem.) under organic and inorganic system of culture. **Ph.D. Thesis**, Tamil Nadu Agric. Univ., Coimbatore.
- Chavan, P., J. Syedismaiz, G.B. Rudraksha, G. Malewar and M.I. Baig. 1997. Effect of various nitrogen levels through FYM and urea on yield, uptake of nutrients and ascorbic acid content in chilli (*Capsicum annum* L.). **J. Indian Soc. Soil Sci.**, **45**: 833-835.
- Deva Kumar N., G.G.E. Rao and S. Shuba.2011. Evaluation of locally available media for the growth and development of nitrogen fixing micro- organisms. **In: Proceedings of the 3rd scientific conference of ISOFAR Organic are life knowledge for tomorrow, held on 28th September-01 October 2011, Korea. PP 504- 509.**
- Elias - Azar, K. 1980. Biocarbonate extractable phosphorus in fresh and composted dairy manure. **Soil Sci. Soc. Am. J.**, **44(2)**: 434-435.
- Goveanthan, A.S., Sugumaran M.P. and Somasundaram, E. (2019). Bio Chemical analysis of Beejamruth and Its Plant Promoting Factors. *Int.J.Curr.Res.Aca.Rev.* 7(5), 1-4.
- Goveanthan A.S., Sugumaran M.P., Ganesh Kumar Gudimetha, Akila S, Suganya K and Somasundaram E (2020). Studies on organic inputs (Jeevamruth and Beejamruth) and their efficacy on fenugreek, *The Pharma Innovation Journal* 9(11), 92-94
- Goveanthan, A.S., Sugumaran, M.P. and Somasundaram, E. (2021). Scientific validation of organic liquid formulation-Jeevamruth by studying its characteristics. *Internat. J. Plant Sci.*, 16 (1): 15-18
- Kuttimani, S. 2004. Response of chilli (*Capsicum annum* L.) genotypes to integrated nutrient management. **M.Sc. Thesis**, University of Agricultural Sciences, Dharwad.
- Maskina, M.S., Yajvinder singh and Bijay singh. 1988. Response of wetland rice to fertilizer N in soil amended with cattle, poultry and pig manure. **Bio wastage**, **26(1)**: 1-8.
- Mathan, K.K., K. Appavu and A. Saravanan. 2000. Effect of organics and irrigation levels on soil physical properties and yield of crops under sorghum - soybean cropping system. **Madras Agric. J.**, **87(1-3)**: 50-53.

- Priyadarsini, J. and P.V.N. Prasad. 2003. Evaluation of nitrogen use efficiency of different rice varieties supplied with organic and inorganic sources of nitrogen. **Andhra Agric. J.**, **50(4)**: 207-210.
- Santos, O.S. Dos and F.T. Nicoloso. 1990. Effects of mineral nitrogen, molybdenum and inoculation with rhizobium on common beans. *Revista do Centro de Ciencias Rurais Universidade Federal de Santa Maria*, 20(1-2): 23-25.
- Shashidhara, G.B. 2000. Integrated nutrient management for chilli (*Capsicum annuum* L.) inalfisols of Northern transition zone of Karnataka. **M.Sc. Thesis**, University of Agricultural Sciences, Dharwad.
- Somasundaram, E. 1991. Studies on the direct and residual effect of applied and intercropped *Sesbania rostrata* on rice. **Asian J. Plant Sciences.**, **6(2)**: 282-287.
- Sreekhantan, L. 1987. Integrated phosphorus management in rice based cropping system. **Ph.D., (Thesis)**, Tamil Nadu Agricultural University, Coimbatore.
- Sugumaran, M.P., Akila.S and Somasundaram.E. 2018. Studies on Analyzing the Shelf Life of Panchagavya with Different Alternatives for Ghee. *International Journal of Agriculture Sciences*, 10(24): 7655-7656
- Sugumaran.M.P, Akila,S and Somasundaram E. (2019). Studies on analysis on biochemical characters of leaf over liquid organic inputs (Panchagavya and Jeevamruth) on Maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*. 8(5), 1794-1797.